

Status, Characterization and Conservation Practices of Local Chicken Ecotypes, Ethiopia

¹Addis Getu, ²Kefyalew Alemayehu ¹Atnaf Alebie

¹Department of Animal Production and Extension, University of Gondar, Faculty of Veterinary Medicine, Ethiopia

²Bahir Dar University, Department of Animal Production and Technology, P.O. Box 21 45, Bahir Dar, Ethiopia

ABSTRACT

Review work was conducted to assess the characterized and conservation methods of indigenous chickens in Ethiopia. In Ethiopia Chickens are the most wide spread and dominant poultry species. Since local chickens have good potential to adapted in different agro-ecology and provide luxurious source of family protein and income to rural poor. However, village chicken is usually kept under free ranging production system. Still those local chickens are non-descriptive type and show variations in body position, color, comb type and productivity. Indigenous chickens have characterized as; poor appearance, relatively low productivity, slow growth rate, small adult size and lays small egg. So, they are neglected from researchers, development workers and policy makers to put them in the research and development. To decrease loose of chicken genetic resource, phenotypic and genotypic characterization work were conducted. However, many chickens are lack with information about their geographical distributions and its availability. Many previous reports underlined that the breed characteristics of indigenous chickens are vary in color, comb type, body conformation and weight. High incidences of chicken diseases, mainly (NCD), coccidiosis, salmonellae's fowl pox are the major and economically important constraint for village chicken production system following feeds and predators. Further constraints are poor access to markets, weak institutions, and lack of skills and knowledge which lead to high rate of genetic erosion. According to DAD-IS and DAGR-IS, the evidences about the genetic resource of identified chickens are undocumented and unobserved as well, only small number of chickens ecotype such as Tilili, Horro, Chefe, Jarso, Tepi, Gelila, Debre-Elias, Melo-Hamusit, Gassay/Farta, Guangua, Mecha, Konso, Mandura, and Sheka are the major identified and characterized type of local chicken ecotypes in Ethiopia. Therefore, conservation practices are not common rather than aggravating the erosion of local chicken resources through random distribution of exotic chickens.

Keywords: Chickens, Conservation, Indigenous

I. INTRODUCTION

In Ethiopia, village chicken production systems usually kept under free range system and their feed is obtained through scavenging. The major feed resource are insects, worms, seeds and plant materials, with very small amounts of grain and table leftover supplements from the household (Tadelle and Ogle, 2000; Bogale, 2008). These scavenging chickens are the most widespread which provide important source of family protein and incomes (Tadelle *et al.*, 2003). At country level local chicken are estimated as 49.3 million (CSA, 2011). Local chickens are non-descriptive type show a large variation in body position, color, comb type and

productivity which also attributed to their widespread distribution and huge population size (Tadelle *et al.*, 2003; Halima, 2007; Fisseha *et al.*, 2010b). Ethiopia is the home of domestic animal migration from Asia to Africa which plaid a great impact to widespread distribution in a country (Halima, 2007). Adaptation of harsh environment and resistance to disease are the major opportunities of local chicken in Ethiopia and contributed to the national economy in general and the rural economy in particular 99.2% of meat and 99% of egg productions are contributed by local chickens with an average annual output of 72,300 and 78,000 metric tons of meat and egg production respectively (Tadelle *et al.*, 2003; Hailemariam *et al.*, 2006; Fessiha *et al.*,

2010b). However, indigenous chickens are poor appearance, relatively low productivity, slow growth rate, small adult size and lays small egg size (Pedersen, 2002; Gondwe, 2004). Due to this effect they are not getting attention by concerned bodies (Tadelle, 2003; Mekonnen, 2007). Therefore the genetic resources in some part of Ethiopia are becoming critically endangered (Halima 2007; Dana *et al* 2010; Dana, 2011). Furthermore, the extensive and random distribution of exotic chicken breeds is cause of dilution in indigenous chickens (Tadelle *et al.*, 2003). To reduce loses of chicken genetic resource, some workers have made phenotypic and genotypic characterization of indigenous chicken in some parts of Ethiopia (Tadelle, 2003; Halima, 2007; Dana *et al.*, 2009; Dana, 2011). Therefore the objective of this paper is to assess the status, characterization and conservation practice of indigenous chicken in Ethiopia and to identify risk of their extinction.

II. METHODS AND MATERIAL

Chicken Population in Ethiopia

Ethiopia is one of African countries with a significant population of chicken and covers about 60% of the total population (Mekonnen et al., 1991).The domesticated poultry species are the part of livestock population. Recently, chickens are estimated to be about 49.3 millions of which 97.3%, 0.38 % and 2.32 % of the total chickens are indigenous, hybrid and exotic respectively (CSA, 2011). This report revealed that chicken includes cocks, cockerels, pullets, laying hens, non-laying hens and chicks.




Figure 1 : Distribution of chicken population by type in Ethiopia

In Ethiopia, the rural farm households do not keep other domesticated birds (Bogale, 2008). The same study indicated that the mean number of breeding females per households was 5.4 ± 2 and the overall male to female ratio of the village flocks was: 1:2.5.

A. Geographical Distributions and Classification of Indigenous Chickens in Ethiopia

There is no complete information about the geographical distributions of many chickens in the developing countries of the world thus some of them are commonly referred to as non-descript breeds (FAO, 2011). Ethiopian local chickens are none descriptive (none have common defined name). As a result breed characteristics of local chickens are vary in color, comb type, body conformation and body weight (Alemu & Tadelle, 1997; Meseret, 2010). The diversity of Ethiopian local chicken shows variation in morphology, color etc and their name is given based on name of place where they are found. Based on their color, the local chicken ecotype classified and named as: *Tukor* (black), *Melata* (nacked nack), *Kei* (red), *Gebsuma* (mixed) and *Netch* (white). Based on feather morphology local chicken ecotypes are categorized as skin color (*silky, white & yellow*), Comp type (*single, rose, pea, walnut & duplex*) and Body shape (*blocky, triangular & wedge*) (Dana *et al.*, 2010). Therefore Dana (2011) currently reported that local chickens are characterized by pronounced broodiness (maternal instinct), slow growth rate, late sexual maturity and low production performance.

B. Importance of Poultry Productions

Word poultry refers to all domesticated birds that are reared for the production of meat and eggs for human consumption as well as for economic benefits. Village chicken production systems are characterized by low input- low output levels and free range production systems (Bogale, 2008). Indigenous chicken provides relevant contributions for poverty alleviation by adapting different agro - ecology, resistance to disease and supplying high quality protein to balance the family food supply, and provide small disposable cash income in addition to the socio-religious functions that are important in the rural people's lives under lack of supplementary feed and other managerial problems. Due to the importance of local chicken, characterizing

and conservation of indigenous chicken genetic resource are mandatory (Tadelle, *et al.*, 2003b). Chicken production is an appropriate and locally available resource on livestock populations. In Africa chicken population is the highest and every rural poor participated in an extensive production system (Tadelle *et al.*, 2003b). From sub-Saharan Africa, 85% of all households keep chicken under free range/ extensive production system, with women owning 70% of it, providing insufficient animal protein in the form of meat and eggs as well as reliable source of cash income (Sonaiya and Swan, 2004; Abubakar *et al.*, 2007). Ethiopia is one of the few African countries which have large chicken population of 60% of the total free range population in Africa (Mekonnen *et al.*, 1991). However, the number of chicken flocks per household in most Ethiopian rural communities is small constituting an average of 7–10 mature chicken, 2–4 adult hens, a male bird (cock) and a number of growers of various ages (Tadelle and Ogle, 2000). Traditional production system of local chickens is common and characterized by their low input and low output levels no feed supplementation, small flock size and periodic devastation of the size by disease and other constraints (Tadelle and Alemu, 1997; Meseret, 2010).

C. Performance of Indigenous Chicken

Some workers reported that indigenous chickens are poor appearance, low performance and produce small sized eggs, slow growth rate, late maturity and slow age at first mating, small clutch size, and high mortality of chicks (Bogale, 2008; Fisseha, 2009; Meseret, 2010). In different part of Ethiopia, comprehensive report stated that phenotypic performance of local chickens are varied because of genotype, management and seasons, such as egg production performance; described in 30 to 60 eggs/hen/yr (Kidane, 1980) at WADU, 34 eggs /hen/yr (Pearson and Brannang, 1990) at Asella, 18-57 eggs/year/ hen (Halima, 2007) at Northwest Ethiopia, 20-60 eggs/3clutch/yr (Bogale, 2008) at Fogera, 10.05 ± 0.15 egg/clutch 3.78 ± 0.07 /yr (Fissiha, 2009) at Bure and the other recent study reported local chicken eggs laid ranges from 53-60 egg/hen/yr range of 43.2-46.96 egg weight (Fisseha *et al.*, 2010a) at North-west Ethiopia.

D. Major Constraints of Poultry Production in Ethiopia

High incidence of chicken diseases, mainly (NCD) is the first and economically important constraint for village chicken production system following by feeds (Dana and Ogle, 2000; Halima, 2007). The other comprehensive study showed that (NCD) is highly infectious and more losses than any other diseases in the tropics and it spreads rapidly through the flock and mortality could reach up to 100% (Dana *et al.*, 2003; Serkalem *et al.*, 2005; Nwanta *et al.*, 2008). Among infectious diseases, salmonellosis, coccidiosis and fowl pox are also considered to be the most important causes of mortality in local chicken while predators are an additional causes of loss (Eshetu *et al.*, 2001). According to Tadelle and Ogle (2000), high mortality of chicks under village chicken production in the central highlands of Ethiopia is due to diseases, parasites, predation, lack of feed, poor housing and insufficient water supply. Further village poultry production is constrained by poor access to markets, goods and services, weak institutions, and lack of skills, knowledge and appropriate technologies (Gueye, 2003). Besbes (2009) reported that poor nutrition and health problems are the main constraints. In addition to the above extensive and random introduction of exotic breeds before appropriate characterization, utilization and conservation of indigenous genetic resources is believed to be the main cause of the loss of indigenous resource (Halima, 2007; Besbes, 2009).

E. Phenotypic Characterization of Indigenous Chicken

Phenotypic characterization is affected by agro-climates, ethnic groups, socio- economic, religious and cultural influences in the nature of the qualitative and quantitative traits variation (Halima, 2007). Ethiopia is gateways of domestic animals migration from Asia to Africa and it has further impact on the diversity of Ethiopian chickens (Hallima, 2007). According to FAO (2011) stated that diversified chicken characterization is identifying distinct Animal Genetic Resource /AnGR/ and describing their uniqueness in their environment within specific location and describes any measurable (quantitative trait), adaptable and observable (qualitative) nature of AnGR and evaluate effective population size

and evaluates status their risks (FAO, 2011). Different report stated that indigenous chickens are characterized in different parts of Ethiopia; Teketel (1986) at Awassa/Sidamo; Bogale (2008) at Fogera District (based on plumage colors as, white, red, black, grayish, brown, white brownish, black brownish, and red brownish), Fissiha (2009) at Bure (characterized based on their phenotypic variations in terms of plumage color, shank length, comb type and growth performances). Based on location Tadelles (2003) at Tilili, Horro, Chefe, Jarso and Tepi, Halima (2007) at Tilili, Gelila, Debre-Elias, Melo-Hamusit, Gassay/Farta, Guangua and Mecha and Dana (2011) characterized at Farta, Konso, Mandura, Horro and Sheka. However, only 5 chickens are listed in DAD-IS (FAO, 2008) and 10 in DAGR-IS (DAGRIS, 2008) including those listed in DAD-IS. This small number represented in the databases indicates that locally adapted populations are still un documented (Dana, 2011).

F. Molecular Characterization

Molecular characterization is the major options for breed definition, especially populations which are not well defined to identify unique alleles. DNA-based methods are independent of environmental factors and provide useful information about genetic diversity to supported the global management of genetic resources (FAO, 2007a; 2011) and genetic conservation (Mendelsohn, 2003; Shrestha, 2004). Characterization and conservation of AnGR is important for designing sustainable poverty alleviation (Toro *et al.*, 2006). At molecular level Tadelles, Halima and Dana has been characterized the Ethiopian chickens. Livestock species are sequenced in genome prepares (Burt, 2005) and therefore provides a vast number of microsatellite markers for diversity studies. According to Halima, twenty two microsatellite markers was used based on the degree of polymorphism and genome coverage for the measurement of chicken diversity (Halima, 2007), and application in diversity studies and detailed information in North West Ethiopia. Information is not available on the genetic diversity of Ethiopian local chickens which are important to design effective selection and conservation strategies (Halima, 2007).

Molecular genetic characterization explores polymorphism (magnify and show different form) in

selected protein molecules and DNA markers to measure genetic variation at the population level. Because of the low level of polymorphism observed in proteins and hence limited applicability in diversity studies, DNA-level polymorphisms are the markers of choice for molecular genetics characterization. Assessment of Genetic variability at the DNA level, different classes of molecular markers have been employed to study genetic diversity in chickens such as restriction fragment length polymorphisms (RFLP), random amplified polymorphic DNA (RAPD) markers, amplified fragment length polymorphisms (AFLP), mitochondrial DNA (mtDNA) markers, two types of variable number of tandem repeat (VNTR) loci (mini satellites and microsatellites), and more recently single nucleotide polymorphism (SNP) markers (Weigend and Romanov, 2001), the assessment of DNA marker polymorphism suggests that variability in DNA is a powerful tool for examining diversity within and among individuals, families and populations. In general identification and characterization of animal genetic resources requires information on their population, adaptation to a specific environment, possession of traits of current or future value and socio-cultural importance, which are crucial inputs to decisions on proper utilization and conservation of AnGR (Dana, 2011).

III. RESULT AND DISCUSSION

Conservation of Poultry Genetic Resources

Global management of genetic resources at international level is mandatory (Gandini & Oldenbroek, 1999). Communication and information system or Domestic Animal Diversity Information System (DAD-IS) is being developed by FAO and USID during 1992, with the objective of assisting countries by providing extensive searchable databases and guidelines for better characterization, utilization and conservation of chicken genetic resources (Halima,2007). Such programmes are important because the AnGR have been faced to genetic dilution due to exotic germplasm use, changes in production systems, markets preferences, natural catastrophes, unstable policies from public and private sectors and the availability of very limited funds for conservation activities (Rege & Gibson, 2003; Halima, 2007; Dana, 2011). It should also include the population size of the animal genetic resources, its physical

description, adaptations, uses, prevalent breeding systems, population trends, predominant production systems, description of the environment in which it is predominantly found, indications of performance levels (meat, growth, reproduction, egg) and the genetic distinctiveness of the animal (Weigend & Romanov, 2002). This provides a basis for distinguishing among different animal genetic resources and for assessing the available diversity (FAO, 2011).

However, insufficient attention has been given to evaluating these resources or to setting up realistic and optimum breeding goals for their improvement (Dana, 2011). As a result some of the animal genetic resources of Africa are endangered, and unless urgent efforts are taken to characterize and conserve, they may be lost even before they are described and documented and it is also stated that an increasing loss of genetic diversity has been observed and poultry genetic resources are considered to be the most endangered (Crawford, 1990; Halima, 2007). The majority of livestock genetic diversity is found in the developing world where

documentation is scarce and risk of extinction is highest and increasing. More particularly, it is estimated that 35 % of mammalian breeds and 63 % of avian breeds are at risk of extinction. These local chickens face genetic erosion which may lead to the loss of valuable genetic variability in specific characteristics of their unique genes and alleles pertinent to their adaptation to particular environments (Romanov *et al.*, 1996). More over characterization, conservation and use of indigenous animal resources under low levels of input in the tropics are usually more productive than is the case with exotic breeds (Halima, 2007).

Further the rule and strategy of the risk indicators of AnGR guide line established for the next conservation measures of AnGR (FAO, 2007a; 2011). So far status of animal Genetic Resource (AnGR) are classified as critical, critical-maintained, endangered, endangered-maintained, extinct, not at risk and unknown breeds (UNEP, 2008). The important points are relevant to show indicators of the risk of the breeds are described in table 1.

Table1. Classification of AnGR based on their endangerments

Groups of breeds	Total number of breeding		Overall population size	Remark
	Male	Female		
Extinct	0	0	0	There is no breeding male and female
Critical	≤5	≤100	≤120	
Critical-maintained				Active conservation programmes
Endangered	5≤20	>100 ≤1000	>1000 or ≤1200	
Endangered-M				Active conservation programmes
Not at risk				Identified and characterized
Unknown				Not identify and characterize

Source: (FAO, 2007a; UNEP, 2008)

The above indicator is an important for sustainable management of genetic resources in respect to genetic conservation measures. Cryopreservation is an important complementary measure for the conservation of diversity in poultry as in other farm animal species. Some recent papers summarize the state of the art in long-term storage techniques for avian semen (Blesbois and Labbè, 2003). Over the past 50 years, preservation technologies have been developed for mammalian

gametes and embryos, in particular in cattle, which enable to run programs to preserve genetic materials (Gibons et al, 2006).

Opportunities of Characterization and Conservation of Chicken Genetic Resource

Since 2005, massively parallel DNA primer sequencing has become available and has reduced the cost of DNA sequencing (Shendure and Ji, 2008). In the next few years, even more effective sequencing systems will be

accessible. This development will open the door for very low cost including the whole genome sequencing. The focus is now moving from the sequencing of a single individual to hundreds or even thousands of individuals. The “1000genome project” in humans which aims to sequence the genomes of approximately 1200 individuals from 3 major populations at approximately increase 4x coverage that targets 10,000 vertebrate species’ Affordable re sequencing will largely extend SNP identification. Analysis of breeds that have not been subject to selective breeding will forward the compilation of bias-free SNP panels for diversity studies. The widespread existence and importance of Copy Number Variation (CNV) on the level of gene expression (Beckman et al., 2007) and of micro RNAs on gene regulation (Shivdasani, 2006) are recent examples of unexpected discoveries. High sequencing in put will be more and more integrated with new statistical approaches (Luikart *et al.*, 2003 ;). We expect that these new tools will stimulate the identification and understanding of variation supporting important traits, including phenotypes relevant for adaptation and sustainable operation. With regard to conservation, whole-genome sequencing will also provide more objective indications of uniqueness than any marker panel (FAO, 2007a). In addition, adaptive variation will be included in prioritization protocols in order to ensure conservation of unique adaptive variants, thus optimizing conservation efforts both in vivo and in vitro. It is also envisaged that breeding and selection will be more and more guided by molecular analysis. Models are to be developed and customized to populations with different genetic structure (small vs large breeds) and to different purposes (genetic improvement, control of inbreeding, maintenance of diversity).

IV. CONCLUSION

The local chicken genetic resource in Ethiopia is play significant role in poverty alleviation generation additional income and religion or cultural reason. Furthermore some worker tried to phenotypic characterization of local chicken in some parts of Ethiopia such as Tillili Tepi Medura Sheka Horo, Jerso, Farta etc. and only 10 types are recognized and documented in DAD-IS and DAGR-IS. Here in African 60% of wide spread distribution and large population size chicken population are found in Ethiopia. However,

this identified small number represented in the databases indicates the shortage of data on chicken genetic resources of Ethiopia suggesting that much of the diversity that exists in the locally adapted populations still remains undocumented. Even if the above limited effort and small number of identified ecotype still didn’t include chicken genetic resource of all parts of Ethiopia like northern Amahara region of north Gonder administrative zone. The indigenous chicken genetic resource of north Gonder zone needs intensive identification, characterization properly utilization and conservation based on their risk status of extinction.

Therefore based on this paper my research is highly initiated to identify and characterize distinct local chicken ecotypes in terms of physical characteristics and production systems of north western part of Ethiopia.

V. RECOMMENDATION

- ✓ As a result, there is a need to design and implement a research programme to collect, conserve and improve the indigenous chickens.
- ✓ Phenotypic characterization should be supported by genetic characterization of animal genetic resource.
- ✓ Lack of documentation, proper utilization and conservation of local Chicken genetic resource is observed.
- ✓ Researchers, developmental workers and policy makers should be give infancies for local genetic resource.
- ✓ Concerned bodies should give mediate response for conservation measure after characterization and describing the risk level of local chicken in Ethiopia.

VI. REFERENCES

- [1] Abdelqader, A., Wollny, C. and Gauly, M., 2007. Characterization of local chicken production system and potential under different level of management practice in Jordan. *Journal of Tropical Animal Health and Production.* 39: 55–164.
- [2] Akishinonomiya, F., Miyake, T., Takada, M., Shingu, R., Endo, T., Gojobori, T., Kondo, N. and Ohno, S., 1996. Monophyletic origin and unique dispersal patterns of domestic fowls. *Proceedings of the National Academy of Sciences of the United States of America.* 93: 6792- 795.
- [3] Alemu Yami, 1995. Poultry production in Ethiopia. *World’s Poultry Science Journal.*51: 197- 201.

- [4] Beckmann, J.S., Estivill, X., Antonarakis, S.E., 2007. *Nat. Rev. Genet.* 8:639-646.
- [5] Besbes, B., 2009. Genotype evaluation and breeding of poultry for performance under suboptimal village conditions. *Food and Agriculture Organization of the United Nations, World's Poultry Science.* 65: 260-271.
- [6] Blesbois, E. and Labbe, C., 2003. Main improvements in semen and embryo preservation for Fish and fowl. *Workshop on Cryopreservation of Animal Genetic Resources in Europe, Paris, France.* Pp. 55-57.
- [7] Bogale Kibret, 2008. In situ characterization of local chicken eco-type for functional traits and production system in Fogera District, Amhara regional state; M.Sc. Thesis Submitted to the Department of Animal Science School of Graduate Studies, Haramaya University.
- [8] Brannang, E. and Person, S., 1990. *Ethiopian animal husbandry and Breeding in the Tropics and Sub-tropics.* Humboldt University of Berlin, Germany. Uppsala, Sweden. 127p.
- [9] Burt, D.W., 2005. Chicken genome: Current status and future opportunities. 15:1692-1698.
- [10] Crawford, R.D., 1990. Origin and history of poultry species. In: *Poultry Breeding and Genetics* (Ed.by R.D. Crawford), Elsevier Science Publishers, Amsterdam, the Netherlands. Pp.1-41.
- [11] CSA, 2007. Population and Housing Census of Ethiopia: Statistical Report for Amhara Region. Pp. 1-25: In Part I: Population Size and Characteristics.
- [12] CSA, 2011. Agricultural sample survey 2010/11. volume 2: statistical bulletin 505. Report on livestock and livestock characteristics (prevent peasant holdings), Addis Ababa, February 2011.21
- [13] DAGRIS, 2008. Domestic Animal Genetic Resources Information System (DAGRIS). International livestock research institute, Addis Ababa, Ethiopia. (<http://dagris.ilri.cgiar.org>). (Accessed on 28 September, 2008).
- [14] Dana N, 2011. Breeding programs for indigenous chicken in Ethiopia Analysis of diversity in production systems and chicken populations; PhD. Thesis submitted in fulfillment of the requirements for the degree of doctor at Wageningen University Netherlands.
- [15] Dana N, Alemu Y, Tadelle D, and Samuel W/H, 2003. On-station and on-farm evaluation of the hay-Box chick brooder using different insulation materials at Debre Zeit Agricultural Research Center and Adaa woreda. Proceedings of the 10th annual conference of the Ethiopian Society of Animal Production (ESAP), August 21–23, held in Addis Ababa Ethiopia. Pp. 211–213.
- [16] Dana N, Tadelle D, Liesbeth, H., v. and Johan, A., M., 2009. Morphological features of indigenous chicken populations of Ethiopia. *Animal Breeding and Genomics Center, Wageningen University .Animal Genetic Resources,* 2010. 46:11–23.
- [17] Dana N, vander, W., E. and Johan, A., M., 2010. Genetic and phenotypic parameter estimates for body weights and egg production in Horro chicken of Ethiopia. Submitted to *Trop Anim. Health Prod, Animal Breeding and Genomics Centre, Wageningen University Netherlands* 14 July 2010.
- [18] Eshetu Y, Mulualem, E., Ibrahim, H., Berhanu, A. and Aberra, K., 2001. Study of gastro-intestinal helminths of scavenging chickens in four rural districts of Amhara region, Ethiopia. *Rev. Sci. tech off. Int. Epiz.* 20(3): 791–796.
- [19] FAO, 2007a. *Global Plan of Action for Animal Genetics Resources conservation, Rome, Italy.* Pp: 125-128.
- [20] FAO, 2008. Domestic Animal Diversity–Information System (DAD-IS). [<http://www.fao.org/dad-is>]. The Food and Agriculture Organization of the United Nation (FAO). Rome, Italy (accessed on December 4, 2008).
- [21] FAO, 2011. Draft guidelines on phenotypic characterization of Animal genetic Resource. Commission on Genetic Resources for Food and Agriculture Rome. 18-22 July, 2011. 6p.
- [22] Fisseha M, 2009. Studies on production and marketing system of local chicken ecotypes in Bure Wereda, North west Amhara. M.Sc. Thesis, Hawassa University, Hawassa, Ethiopia.
- [23] Fisseha M, Abera M, and Tadelle D, 2010a. Assessment of village chicken production system and evaluation of the productive and reproductive performance local chicken ecotype in Bure district, North West Ethiopia. *African Journal of Agricultural Research* Vol. 5(13): 1739-1748, 4 July, 2010.
- [24] Fisseha M, Azage T, and Tadelle D, 2010b. Indigenous chicken production and marketing systems in Ethiopia: Characteristics and opportunities for market-oriented development. Working paper No.24; Improving Productivity and Market Success (IPMS) of Ethiopian Farmers Project, International Livestock Research Institute (ILRI). Addis Ababa, Ethiopia.66pp.
- [25] Gandini, G.C. & Oldenbroek, J.K., 1999. Choosing the conservation strategy. In: *Genebanks and the conservation of farm animal genetic resources.* Ed. J.K. Oldenbroek. DLO Institute for Animal Science and Health, the Netherlands.
- [26] Gibons, J., Gamange, S., Hanotte, Iñiguez L., Maillard J.C., Rischkowsky B., Semanbo D. and Toll., 2006. "Options and Strategies for the Conservation of Farm Animal Genetic Resources", Report of an International Workshop (7-10 November 2005, Montpellier, France), Biodiversity International, and Rome, Italy: 53 pp.
- [27] Gondwe, T.N.P., 2004. Characterization of local chicken in low input-low output production systems: is there scope for appropriate production and breeding strategies in Malawi? PhD thesis, Georg- ugst- niversität Göttingen, Germany.
- [28] Gueye, E., 2003. Poverty alleviation, food security and the well-being of the human population through family poultry in low income food-deficit countries. *Senegalese Institute of Agricultural research (ISRA), B.P.2057 and Dakar-hann, Senegal.*
- [29] Hailemariam T, Legesse D, Alemu Y, and Dana N, 2006. Adaptation of poultry breeds in the high lands of Ethiopia. *Ethiopian institute of agricultural research.* 26p.
- [30] Halima H, 2007. Phenotypic and genetic characterization of indigenous chicken populations in Northwest Ethiopia. PhD. Thesis submitted to the faculty of National and agricultural sciences department

- of animal Wild life and Grass land Sciences University of the Free State, Bloemfontein, and South Africa.
- [31] Halima, H., Nesor, F.W.C., Tadelles, D., Kock, A. De, Van Marle-Köster, E., 2006b. Studies on the growth performance of indigenous chicken ecotypes and RIR chicken under improved management system in North west Ethiopia. *Livest. Res. Rural Dev.* 18 (6), (<http://www.cipav.org.co/lrrd/lrrd18/6/hass18076.htm>).
- [32] Halima, H., Nesor, F.W.C., Van Marle-Köster, E., Kock, A.De, 2007b. Phenotypic variation of indigenous chicken populations in northwest ethiopia. *trop. anim. hlth prod.* (in press).
- [33] Halima, H., Nesor, F.W.C., Van Marle-Köster, E., Kock, A.De, 2007a. Village-based indigenous chicken production systems in north-west Ethiopia. *Trop. Anim. Hlth Prod.* 39 (3):189-197.
- [34] Kidane, H., 1980. Performance of F1 cross breeds. Welaita Agricultural Development Unit. Animal husbandry and breeding. Welaita Sodo, Ethiopia. Bulletin No. 4. 33p.
- [35] Luikart, G., England, P.R., Tallmon, D., 2003. *Nat. Rev. Genet.* 4:981-994.
- [36] Mekonnen G/r, 2007. Characterization of smallholder poultry production and marketing system of dale, Wonsho and Loka Abaya Weredas of southern Ethiopia. M.Sc Thesis, Awassa College of Agriculture, Hawassa University.
- [37] Mekonnen G/r, Teketel, F., Alemu, G., Dagnatchew, Z. and Anteneh, A., 1991. The Ethiopian livestock industry: retrospect and prospects. Proc.3rd National Livest. Improvement Conference, Institute of Agricultural Research, Addis Ababa, Ethiopia.
- [38] Mendelsohn, R., 2003. The challenge of conserving indigenous domesticated animals. *Ecological Economics.* 45(3):501-510.
- [39] Meseret M, 2010. Characterization of Village Chicken Production and Marketing System in; M.Sc thesis Submitted to the Department of Animal Science, Jimma University, College of Agriculture and Veterinary Medicine, School of Graduate Studies.
- [40] Nwanta, J.A., Egege, SC., Alli-Balogun, JK. and Ezema, WS., 2008. Evaluation of prevalence and seasonality of Newcastle disease in chicken in Kaduna, Nigeria. *World's Poultry Science Journal.* 64:414-416.
- [41] Pedersen, C.V., 2002. Production of semi-scavenging chicken in Zimbabwe. PhD. Thesis, Royal Veterinary and Agricultural University, Copenhagen.
- [42] Rege, J. E. O. and Gibson, J. P., 2003. Animal genetic resources and economic development: issues in relation to economic valuation. *Ecological Economics* 45 (3): 319-330.
- [43] Romanov, M. N., Wezyk, S., Cywa-Benko, K. and Sakhatsky, N.I., 1996. Poultry genetic resources in the countries of Eastern Europe: History and current state. *Poult. Avian Biology Rev.* 7: 1-29.
- [44] Serkalem Tadesse, Hagos Ashenafi, and Zeleke Aschalew, 2005. Sero-prevalence study of Newcastle disease in local chickens in central Ethiopia. *International Journal of Applied Research. Vet. Med.* Vol. 3, No. 1/19.
- [45] Shivdasani, R.A. 2006. Blood Sampling. *108:3646-3653.*
- [46] Shrestha, J.N.B., 2004. Conserving domestic animal diversity among composite populations. *Small Ruminant Res.* 56(1-3): 3-20.
- [47] Sonaiya, E.B. and Swan, S.E., 2004. Small-scale poultry production, technical guide manual. FAO Animal Production and Health 1. FAO (Food and Agriculture Organization of the United Nations), Rome, Italy.
- [48] Tadele D, 2003. Phenotypic and genetic characterization of chicken ecotypes in Ethiopia. PhD. Thesis submitted to Humboldt University, Germany.
- [49] Tadelles D, Alemu Y and Peters, K., 2003b. Village chicken production systems in Ethiopia: Use patterns and performance valuation and chicken products and socio-economic functions of chicken. *Livestock Research for Rural Development* 1(15). (Available from <http://www.lrrd.org/lrrd15/1/tadeb151.htm>) (Accessed on 1 September 2010).
- [50] Tadelles D and Ogle, B., 2000. Nutritional Status of Village Poultry in the Central Highlands of Ethiopia as Assessed by Analyses of Crop Contents. Department of Animal Nutrition and Management, Debre Zeit Agriculture Research Centre, Debre Zeit, Ethiopia. *J.Agric.Sci.* 17: 47-56.
- [51] Toro, M.A., Fernandez, J. and Caballero, A., 2006. Scientific basis for policies in conservation of farm animal genetic resources. Proc. 8th World Congr. Genet. Appl. Livest. Prod., Belo Horizonte, MG, Brasil.
- [52] UNEP, 2008. Consideration of an intergovernmental science-policy platform on biodiversity and ecosystem services: objectives and functions of an intergovernmental science-policy platform on biodiversity and ecosystem services. Kuala Lumpur, 10-12 November 2008. 3P.
- [53] Weigend, S. & Romanov, M.N., 2002. The World watch list for domestic animal diversity in the context of conservation and utilization of poultry biodiversity. *World's Poult. Sci.* 58 (4): 411- 430.
- [54] West, B. and Zhou, B., 1998. Did chickens go north? New evidence for domestication. *Journal of Archaeological Science.* 15: 515-33.