

pH-Metric Study on Metal-Ligand Stability Constants of Co(II), Cu(II), Ni(II) and Zn(II) Complexes with some Substituted Novel Chalcones

P. J. Parmar

Department of Chemistry, Nanasaheb Y. N. Chavan Arts, Science and Commerce College, Chalisgaon, Jalgaon, Maharashtra, India

ABSTRACT

The interactions of Co(II), Cu(II) and Zn(II) with 1-(2"-Hydroxy-3"-nitro-5"-methylphenyl)-3-phenyl-2-propen-1-one (Ligand-I) and 1-(2"-Hydroxy-3"-nitro-5"-methylphenyl)-3-(4'-methoxyphenyl)-2-propen-1-one (Ligand-II) have been studied by Bjerrum method as adopted by Calvin and Wilson. The stability constant of 1:1 and 1:2 complexes of Co(II), Cu(II) and Zn(II) have been studied at constant temperature ($27 \pm 0.1^\circ\text{C}$) and 0.1 M ionic strength (NaOH) in 70% dioxane-water mixture. It is observed that formation of 1:1 and 1:2 complexes in occurring simultaneously.

Keywords : pH-Metric Study, Co(II), Cu(II), Ni(II) and Zn(II), Metal-Ligand Stability Constant, Substituted Novel Chalcones.

I. INTRODUCTION

Chalcones constitute an important group of natural products and some of them possess wide range of biological activities such as antibacterial^{1,2}, anticancer^{3,4}, antitubercular⁵, antiviral^{6,7}, antiinflammatory⁸, etc. The presence of a reactive, α,β -unsaturated keto function in chalcones is found to be responsible for their antimicrobial activity, which may be altered depending on the type and position of substituent on the aromatic rings. In the present communication, the reaction of 2-hydroxy-3-nitro-5-methyl acetanaphthone (1) with different aromatic aldehydes (2) to form chalcones (Ligand I and Ligand II) is reported. The structures of the various synthesized compounds were assigned on the basis of elemental analysis, IR and ^1H NMR spectral data. These Compounds were also screened for their antimicrobial activity.

Since the last four decades considerable research work has been done on the study of complexes in solution^{9,10}. Bjerrum's dissertation¹¹ has taken the

initiative to develop the field. Some of the important characteristics added to the drugs play an important role in biological activities, e.g., low dissociation constant, special redox potential, electron distribution and solubilities which has been imparted by the metal complex formation. The effect on solubilities of the drug in the lipids and their transfer through the cell membrane have been marked by these characteristics. The group transfer reactions, bond formation, bond cleavage are the natural processes which have been processed by metal-complex formation. Metal complexations not only bring the reacting molecules together to give activated complex¹² but also polarize electrons from the ligands towards the metal. The relation between stability and basicity of the ligands is indicated by the formation constant and free energy charge value. Bulkier group increases the basicity of the ligands as well as stability. The stability of the complexes is determined by the nature of central metal atom and the ligands. The stability of the complex compounds is influenced by the most important characteristics like the degree of oxidation, the radius and electronic structure. Stability of the


The ionic strength of all the solutions was maintained constant (0.1 M) by adding appropriate quantity of 1 M KNO₃ solution.

The titrations were carried out in 100 mL pyrex glass beaker kept in a ice-cold water bath maintained at constant temperature (27±0.1°C). Nitrogen gas was purged for chemically inert atmosphere. The readings were recorded for each addition of 0.5 mL. The graph of volume of alkali against pH were plotted. The pH-meter readings were taken only after the gas bubbling completely stopped.

III. RESULTS AND DISCUSSION

Determination of proton-ligand stability constants

The dissociation constants of substituted diketones (1, 2) were determined at 0.1 M ionic strength pH-metrically. The reagents or ligands used in the present investigation are monobasic acids having only one dissociate H⁺ ion from OH group and can therefore be represented as HL.



The titration curves of the acid and the ligand deviate at about pH 1.5 and then increase upto pH 7.5. The deviations between an acid curve from ligand for all the systems showed the dissociation of H⁺ ions from –OH groups of the ligands.

Proton-ligand formation numbers (η_A) were calculated by the Irving and Rossotti expression.

$$\eta_A = Y \frac{(E^{\circ} + N) \Delta V}{(V_0 + V^1) T^{\circ} L}$$

Where V₀=Initial volume of solution(50 mL)

N=Normality of sodium hydroxide

T[°]L =Normality of Sodium hydroxide

E[°]=Initial concentration of free acid(HNO₃).

Y=Number of dissociable protons from ligand.

(V₂-V₁)=Δ V=Volumes of alkali consumed by acid and ligand

On the same pH

The pK_a values for both systems were calculated by half integral calculation which are presented in Table 1.

Table 1. Proton-Ligand Stability Constants

System	pK
Ligand-1: 1-(2"-Hydroxy-3"-nitro-5"-methylphenyl)-3-phenyl-2-propen-1-one	6.97
Ligand-2: 1-(2"-Hydroxy-3"-nitro-5"-methylphenyl)-3-(4'-methoxyphenyl)-2-propen-1-one	6.69

Table 2. Metal-Ligand Stability Constants(Log K)

System	Metal-ligand stability constant	
	log K ₁	Log K ₂
Ligand-1 Co(II)	8.15	6.71
Ligand-1 Cu(II)	8.83	6.45
Ligand-1 Zn(II)	8.57	6.55
Ligand-1 Ni(II)	8.41	6.29
Ligand-II Co(II)	7.35	5.95
Ligand-II Cu(II)	6.75	5.57
Ligand-II Zn(II)	8.27	6.35
Ligand-II Ni(II)	8.11	6.17

IV. CONCLUSION

It is observed from the titration curves between (acid+ligand) and (acid+ligand+metal) for all the systems that the deviation between the curves is found to be at about pH 1.5. This indicates the commencement of complex formation from this pH. During the titration process, solution colour is changed from light to brown to dark brown. This also shows the formation of complex between ligand and metal ion.

It could also be seen from Table 1 that the order of pK values of the ligands is found to be as:

pK ligand 1 > pK ligand 2 this may be due to the fact of the presence of nitro(-NO₂) as the electron withdrawing groups in ligand 2 in addition to methoxy group (-OCH₃) as the electron donating group at 4-position of second phenyl ring in the same ligand 2. The electron withdrawing group reduces the pK values of the ligand.

It is observed from Table 2 that the difference between the values of log K₁ and log k₂ is sufficiently large. This indicates the stepwise complex formation between ligand and metal ions. The values of metal-ligand stability constants (log K₁ and log K₂) are greater of Cu(II) complexes than Co(II) and Zn(II) complexes. It means Cu(II) forms more stable complexes than the complexes of Co(II) and Zn(II) with ligand 1 but less stable in case of ligand 2.

V. ACKNOWLEDGEMENT

Thanks to Principal Nanasaheb Y. N. Chavan Arts, science and Commerce College, Chalisgaon for providing necessary laboratory facilities and valuable support for conducting this research work.

VI. REFERENCES

- [1]. S. Ishidia, A. Mabsuda, Y. Kawamuna and Yamanaka, *Chemotherapy*, 8, 146 (1960).
- [2]. M. B. Hogale, N. P. Dhore, A. F. Shelar and P. K. Pawar, *Orient. J. Chem.*, 2, 55 (1986).
- [3]. J. Matthew, A. V. S. Rao and S. Rambhav, *Curr. Sci.* 53, 576 (1984).
- [4]. T. Yamakawa, H. Kagechika, E. Kawachi, Y. Hashimoto and K. Shedo, *J. Med. Chem.*, 33, 1430 (1990).
- [5]. V. K. Ahluwalia, L. Nayal, N. Kaila, S. Bala and A. K. Tahim, *Indian J. Chem.*, 26B, 384 (1987).
- [6]. A. K. Bhatt, R. P. Bhamana, M. R. Patel, R. A. Bellare and C. V. Deliwala, *Indian J. Chem.*, 10, 694 (1972).
- [7]. H. Ishitsuka, Y. T. Ninomiya, C. Ohsawa, M. Fujju and Y. Suhara, *Antimicrob. Agents Chemother.*, 22, 617 (1982).
- [8]. Y. Ninomiya, N. Shimma and H. Ishitsuka, *Antiviral Res.*, 13, 61 (1990).
- [9]. F. H. C. Rossoti and H. Rosotti, *The determination of stability constants*, McGraw-Hill Book Co.
- [10]. A. E. Martell and M. Calvin, *Chemistry of the Metal Chelate Compounds*, Prentice Hall, Inc., England, Cliffs, N. J. (1962).
- [11]. J. Bjerrum, *Metal Amine Formation in Aqueous Solutions*, P. Haase and Sons, Copenhagen (1941).
- [12]. A. T. Florence and D. Attwood, *Physical Principles of Pharmacy*, Mcmillan, London (1981).
- [13]. H. Arving and R. J. P. Williams, *J. Chem. Soc.*, 3192 (1953).
- [14]. Z. Ley, *Electrochem.*, 10, 954 (1904).
- [15]. Pfeiffer, *Angew Chem.*, 53, 93 (1940).
- [16]. Schwarzenbach and Ankerman, *Helv. Chim. Acta*, 31, 1029 (1948).
- [17]. V. Fukuda, R. Morishita and K. Sone, *Bull Chem. Soc. Jpn*, 49, 1017 (1976).

- [18]. A. K. Rana and J. R. Shah, J. Indian Chem. Soc., 63, 281 (1981).
- [19]. S. Gudadhe, M:L. Narwade and V. S. Jamode, Acta Cienc. Indica(Chem.). 11,234(1985).
- [20]. C. Natrajan and P. Thormaraj, Indian J. Chem., 30A, 722 (1991).
- [21]. V. B. Tayade, Synthesis, Studies of heterocyclic compounds containing O, N and Sbeteroatoms, Ph. D. Thesis, Amravati University, Amravati (1998).
- [22]. Ya. A. Shuter, Zh. Obshch, Khim., 44,379 (1974).
- [23]. J. C. Jansen and J. Reedijk, Z. Naturforsch, Teil B, 29, 527 (1974) (Eng.).
- [24]. V. S. Jolly, G. D. Arora and P. Taiwar, J. Indian Chem. Soc., 67, 1001 (1990).