

Protective Effect of Curcumin Longa on Sodium Arsenite Induced Toxicity in Swiss Albino Mice

Dr. Pramod Shanker

TGT (Science),

S.R.P.S.Govt.Sr.Sec.School,

Gardanibagh, Patna, India



Abstract- Arsenic in the present times has caused serious health hazards in the human population. In India, the arsenic poisoning in ground water in Gangetic plains has increased many folds. The exposed population are suffering from various arsenicosis symptoms such as hyperkeratosis, melanosis, and other internal organ diseases of gastrointestinal, cardiovascular, hormonal and others. Hence, the present study is focused to combat the deleterious effect of arsenic toxicity in animal models utilizing medicinal plant extract. The animals (Swiss albino mice) were treated with Sodium arsenite at the dose of 8 mg per kg body weight for 16 weeks to make arsenic model and upon these arsenic pre-treated rats rhizome extract of Curcuma longa at the dose of 25 mg per kg body weight was administered for 4 weeks to study the ameliorative effects of this plant extract. After the entire treatment, rats were sacrificed and their blood samples were obtained and analysed for haematological and biochemical study. The study shows that arsenic induced toxicity caused deleterious effect on the mice at the haematological and biochemical levels and there was significant normalisation in the animal at all the respective levels. Hence, Curcumin longa possesses ameliorative properties against arsenic induced toxicity and can be used for human purpose after dose titration.

Keywords: Sodium arsenite, Curcumin longa , protective effect, Swiss albinomice

I. INTRODUCTION

Arsenic menace in the recent years has become a major challenge in the terms of its health impact. About more than 300 million people are affected with the arsenic poisoning due to its contamination in ground water. In the state of Bihar about 10 million populations are exposed to arsenic due to drinking of contaminated water causing serious health related problems. The Gangetic plains are the most affected area in the state where the exposed populations along with common disease burden are also suffering from the non-communicable disease like diabetes. The trivalent form of arsenic is more toxic than the pentavalent form and unfortunately, in exposed population groundwater mostly trivalent form has been observed.

It enters the human body through drinking water via human blood and causes various changes at the level of biochemical, hormonal, tissue and gene level disrupting the normal function of the cells. The also cause skin manifestations according to the magnitude of the exposure. The long duration exposure of arsenic causes damage to the vital organs of the body such as Kidney lungs, urinary bladder, liver, hypertension etc. (Hartwig et al., 2002; Andrew et al., 2006). Arsenic usually binds with the sulfhydryl groups of enzymes which in turn causes changes in the epidermal keratinocytes leads to cause skin manifestations such as keratosis, melanosis, rain drop pigmentation etc. (Patlolla et al., 2005).

Use of medicinal plants for the ailments of various diseases has attracted researchers to decipher novel plants against the disease as it has very good impact and very least side effects (Maurya et al., 2011). Plethora of medicinal plants are available as antioxidants but, study on antidote for arsenic induced toxicity is very less. One such potential plant source is rhizome extracts of Curcumin longa . The phytochemical investigations of Curcumin longa revealed that it contained bioactive constituents such as cuminaldehyde, curcuma longa.

Thus, the present study was undertaken to evaluate the antidote and antitoxic potential of the ethonolic extract of curcuma longa on sodium arsenite induced hepatotoxicity and nephrotoxicity in mice.

II. Experimental procedures

Animals:

Adult and healthy Swiss albino mice (24) of 8 weeks old weighing around 28g to 32g were used for the present study which were provided by the animal house of Mahavir Cancer Sanasthan and Research Centre, Patna, India (CPCSEA Reg-No. 1129/bc/07/CPCSEA). Ethical approval was obtained from the Institutional Animal Ethics Committee (IAEC) with IAEC No. IAEC/2012/12/04. The mice were acclimatized to laboratory housing conditions under 12 h light and dark cycles (room temperature maintained at $22 \pm 2^\circ\text{C}$) for 15 days prior to the beginning of the treatment under standard laboratory conditions. These experimental mice were housed in conventional polypropylene cages with stainless steel grill top and the diet including food (self prepared laboratory) and water to mice were provided *ad libitum*.

Chemicals:

Arsenic was used as Sodium Arsenite (98.5%) manufactured by Sigma-Aldrich, USA (CAS Number: 7784-46-5; S7400-100G), Lot# SLBH5736V, PCode 1001683292. It was purchased from the Scientific store of Patna, Bihar, India.

Preparation of *Curcuma longa* ethanolic extract: In the present study *Curcuma longa* rhizome were purchased from local market of Patna, Bihar and identified by Dr. Ramakant Pandey (Botanist),

Department of Biochemistry, Patna University, Patna, Bihar, India. The plant rhizome were washed through running tap water and incubated at 37°C temperature. The dried rhizome were grinded to fine powder which was further soaked in 100% ethanol for 48 hours, and finally extracted using Rota vapour apparatus after continuous drying for 4 hours at pressure 4psi. The ethanolic extract dose was calculated after LD₅₀ estimation and the final dose was titrated to 25 mg/kg body weight. Previously, for LD₅₀ estimation, 8 groups of rats (n=4 each) were set up and different doses of plant extract were orally administered for 7 days. The LD₅₀ dose was found to be 3000mg/Kg body weight.

Experimental Design:

Mice were randomly divided into four groups, each group comprising six mice and categorized as following. Group I: Normal control group. Group II: Arsenic treated- mice were orally sodium arsenite induced (8mg/kg body weight/day) for 16 weeks and were sacrificed after the completion of the experiment. Group III: Arsenic treated control- mice were orally treated with sodium arsenite (8mg/kg body weight/day) for 16 weeks and left without any dosing for further 4 weeks.

IV: *Curcumin longa* extract administration- Mice were pretreated with sodium arsenite (8mg/kg body weight/day) followed by administration of extract of curcumin ethanolic extract (25mg/kg body weight/day) for 4 weeks. After the end of dosing, mice were anaesthetized by diethyl ether and sacrificed. Blood samples were collected through the orbital puncture from all group of mice. Serum were then separated for the various biochemical estimation.

Biochemical assay: Biochemical analysis were performed of the serum by standard kit process (Coral crest) through (UV - Vis) spectrophotometer (UV-10, Thermo Fisher, USA). The Liver Function Test as Serum Glutamic Pyruvate Transaminase (SGPT) and Serum Glutamic Oxaloacetate Transaminase (SGOT) were measured according to the method (Reitman & Frankel, 1957), Alkaline Phosphate (ALP) by method (Kind & King, 1954), total bilirubin activity by method (Jendrassik & Grof, 1983) while albumin level measured by BCG method. The Kidney Function Test (KFT) were analysed through Urea by (Fawcett, 1960 and Berthelot, 1859), Creatinine by (Toro, et al 1975), and Uric acid by (Bones, 1945).

Lipid Peroxidation (LPO):

Thiobarbituric acid reactive substances (TBARS), as a marker of LPO, were evaluated through the double heating method (Draper and Hadley, 1992) based on the principle of spectrophotometric measurement of color reproduced during the reaction to thiobarbituric acid (TBA) with malondialdehyde (MDA). For this study, 2.5 ml of 100gm/L solution of Trichloroacetic acid (TCA) were mixed with 0.5 ml serum in a centrifuge tube and heated in the water bath at 90°C for 15 minutes. After cooling, at room temperature, the mixture was further allowed to centrifuge at 3000 rpm for 10 minutes, and 2 ml of the supernatant was mixed with 1ml of 6.7gm/L TBA solution in a test tube which was further heated in water bath at 90°C for 15 minutes and left for cooling at the room temperature. Thereafter, further absorbance was measured by UV - Visible spectrophotometer (Thermo Scientific UV-10 USA) at 532 nm.

2.7. Statistical analysis:

Results are presented as mean \pm Standard Deviation (SD) for six mice individual groups and total variation represented in a set of data was analyzed through one-way Analysis of Variance (ANOVA). Differences among mean variance has been analyzed by applying Dunnett's 't' test at 99.9% ($p < 0.05$) confidence level. Calculations were performed with the GraphPad Prism Program (GraphPad Software, Inc., San Diego, USA).

III. Results

Haematological Study:

In the present study, the data of haematological parameters are shown in Table.1, and the study shows significant decrease $P < 0.0001$ in the erythrocyte counts (RBCs), haemoglobin percentage, haematocrit percentage, MCV, MCH but significant increase in leukocyte count (WBCs) in comparison with control group after 16 weeks of exposure. But, after administration of *Curcumin longa* extract there was significant reversal in the haematological values. ANOVA showed that the sodium arsenite has more deleterious effect on time duration of exposure ($P < 0.0001$) (Table 1).

Table 1. Changes in the haematological parameters of Swiss albino mice exposed to Sodium arsenite at the dose of 8 mg/Kg body weight daily for 16 weeks and its amelioration *Curcumin longa* extract at the dose of 25 mg/Kg body weight for 4 weeks.

Blood Parameters	Control	Arsenic treated	<i>Curcumin rhizome</i> extract treated
RBC Counts (10^6 /mm ³)	6.753 \pm 0.25	2.98 \pm 0.498	5.32 \pm 0.225
Hb (percentage)	14.22 \pm 0.56	5.33 \pm 1.365	12.1 \pm 0.549
Haematocrit percentage (Hct) (%)	43.10 \pm 1.20	16.0 \pm 2.276	36.02 \pm 0.329
MCV (fL)	63.8 \pm 1.67	53.7.21 \pm 2.13	67.7 \pm 1.51
MCH (pg)	21.1 \pm 0.27	17.9 \pm 3.62	22.7 \pm 1.56
MCHC (g/L)	33.56 \pm 0.036	33.3 \pm 2.12	33.6 \pm 1.45
WBC (10^3 /mm ³)	7240 \pm 3.56	3690 \pm 8.35	8375 \pm 3.90
Platelets (mm ³)	256000 \pm 2.33	45000 \pm 22.10	180000 \pm 2.90

*The data are presented as mean \pm S.D, $n = 6$, significance at $P < 0.0001$.

3.2. Biochemical Assay:

The biochemical assay showed the following findings - in control group of mice, SGPT level was 29.56 \pm 1.231 U/ml and after 16 weeks of arsenic treatment was 202.6 \pm 27.23 U/ml, while, it was

54.21 ± 23.56 U/ml after 4 weeks of *Curcumin longa extract* administration. In control group of mice, SGOT level was 33.51 ± 2.512 U/ml and after 16 weeks of arsenic treatment was 254.1 ± 14.03 U/ml, while, it was 65.22 ± 1.31 U/ml after 4 weeks of *Curcumin longa extract* administration. In control group of mice, ALP level was 7.45 ± 0.539 K.A units and after 16 weeks of arsenic treatment was 35.23 ± 3.421 K.A units, while, it was 12.16 ± 1.09 K.A units after 4 weeks of *Curcumin longa extract* administration. In control group of mice, bilirubin level was 0.853 ± 0.234 mg/dl and after 16 weeks of arsenic treatment was 2.45 ± 0.453 mg/dl, while, it was 1.217 ± 1.20 mg/dl after 4 weeks of *Curcumin longa extract* administration. In control group of mice, the kidney function test urea level was 29.3 ± 1.323 mg/dl and after 16 weeks of arsenic treatment was 86.20 ± 2.189 mg/dl, while, it was 32.18 ± 0.47 mg/dl after 4 weeks of *Curcumin longa extract* administration. In control group of mice, uric acid level was 2.98 ± 0.332 mg/dl and after 16 weeks of arsenic treatment was 13.90 ± 2.343 mg/dl, while, it was 3.89 ± 1.098 mg/dl after 4 weeks of *Curcumin longa extract* administration. In control group of mice, the creatinine level was 0.902 ± 0.198 mg% and after 16 weeks of arsenic treatment was 2.531 ± 1.001 mg%, while, it was 1.209 ± 0.233 mg% after 4 weeks of *Curcumin longa extract* administration. In control group of mice, the lipid peroxidation level was 4.289 ± 0.523 nmol/ml and after 16 weeks of arsenic treatment was 88.23 ± 3.445 nmol/ml, while, it was 11.27 ± 1.224 nmol/ml after 4 weeks of *Curcumin longa extract* administration (Table 2).

Table 2. Changes in the biochemical parameters of Swiss albino mice exposed to Sodium arsenite at the dose of 8mg/Kg body weight for 16 weeks and its amelioration by *Curcumin longa extract* at the dose of 250 mg/Kg body weight for 4 weeks.

Parameters	Control	Arsenic Treated	<i>Curcumin rhizome</i> extract Treated
SGPT (U/mL)	29.56 ± 1.231	202.6 ± 27.23 ***	54.21 ± 23.56 ###
SGOT (U/mL)	33.51 ± 2.512	254.1 ± 14.03***	65.22 ± 1.31 ###
ALP (KA Unit)	7.45 ± 0.539	35.23 ± 3.421 ***	12.16 ± 1.09 ###
Total Bilirubin (mg/dL)	0.853 ± 0.234	2.45 ± 0.453*	1.217 ± 1.20 ##
Urea (mg/dL)	29.3 ± 1.323	86.20 ± 2.189**	32.18 ± 0.47 ###
Uric acid (mg/dL)	2.98 ± 0.332	13.90 ± 2.343**	3.89 ± 1.098 ##
Creatinine (mg/dL)	0.902 ± 0.198	2.531 ± 1.001 **	1.209 ± 0.233 ##
LPO (nmol/ml)	4.289 ± 0.523	88.23 ± 3.445 *	11.27 ± 1.224 #

*# Effect of different treatments on liver and kidney biomarker parameters in the studied groups (n=4, Significant *** $P < 0.0001$, * $P < 0.05$ compared with control group, ### $P < 0.0001$ compared with arsenic treated group, values are expressed as mean ± SEM).

Discussion

In the present study there has been significant changes observed in the studied parameters. There was significant decrease in the RBC indices parameters such as haemoglobin percentage, hematocrit percentage and MCHC as well as in WBC counts and platelets counts. The present result speculates that there has been significant myelosuppression causing the serious failure of the defense mechanism. In the present study there has been significant increase in the lipid peroxidation (ROS) levels. This confirms the breach in the defense mechanism. Moreover, this causes the depletion of lipids from the membrane which is the onset of loss of cellular integrity (Dwivedi et al., 2015; Sankar et al., 2016).

Arsenic usually enters the system through the gastrointestinal tract and interacts with the sulfhydryl moieties and causes disruption in the metabolic functions associated with it (Lu et al., 2007). However, the excess arsenic concentration is eliminated through the urine and bile (Helleday et. al, 2000). But the arsenic absorbed in the liver gets accumulated in the vital organs of the body, interfering the signalling pathways (Soni, et al. 1993). It also interferes with the cellular apoptosis pathways causing abrupted function causing the onset of carcinogenesis of the vital organs such as kidney, bladder, liver etc. (Mallikarjuna et al., 2003; Yamanaka et al. 2004). The methylated DMA⁺⁵ acts as tumour promoter (Yamamoto et.al, 1995 & Wanibuchi, et. al, 1996; Cohen, et al. 2006; Miller et. al., 2002; Huang et al., 2004; Duker et al.,2005).

In the present study there has been remarkable changes observed as there was significant increase in the SGPT, SGOT, ALP, bilirubin, urea, uric acid and creatinine levels. Recent study speculates that the arsenic in the form of Dimethyl arsenic acid (DMA) is converted which is still a carcinogen category I, which disrupts the functions of the liver as well the kidney (Chen et al., 2011; Hsueh et al., 2009; Zheng et al., 2013 & 2014; Jha et al., 2013). The biochemical changes observed could be due to the deleterious effect caused by the DMA.

In the present study, there has been significant restoration due to curcumin extraction administration especially the haematological parameters and the biochemical parameters. Curcumin bears active ingredients like *Curcumin longa*. The major compounds of curcumin are Tetra hydrocurcumin, that exhibits many of the same physiologic and Pharmacological activities as curcumin and in some systems may exert greater antioxidant activity than curcumin. Apart from it also contains phenolic acids, flavonoids which possesses antioxidant activity play a vital role in inhibiting the lipid peroxidation as well as various types of oxidizing enzymes (Li and Jiang 2004; Gallo et al.,2010; Morshedi et al., 2015). However, the other functions of Curcumin has also been studied by various researchers such as antiplatelet (Sekin et al., 2007), antibacterial (Lee 2005), antifungal (Morshedi and Aliakbari), antidiabetic (Bennet et al., 1982), anti-Parkinson's (Johri 2011), while it also activates the salivary glands and facilitates the primary digestion of the food and produce carminative effects. The protective effect of Curcumin longa seed extract has been merely reported. Hence, the active ingredient Curcumin might have played the vital role to control the arsenic induced hepato-renal toxicity, normalizing the metabolic functions of the liver and kidney

Conclusion:

From the entire study it can be concluded that arsenic causes deleterious effect on the studied parameters especially at the haematological and biochemical levels in the mice disrupting the normal functioning of the body. But, after the administration of the Curcumin longa rhizome in them. Hence, it indicates that Curcumin extract possesses hepato-renal protective effect against arsenic induced toxicity. Therefore, it can be targeted as novel and safe antitoxic drug against arsenic.

Declaration of competing interest

The authors declare that they have no conflict of interest.

Acknowledgement

The authors are thankful to Mahavir Cancer Sansthan and Research Centre for providing infrastructural facility during this work.

References :

- [1]. Alizadeh Behbahani, B., Noshad, M., & Falah, F. 2019. Cumin essential oil: Phytochemical analysis, antimicrobial activity and investigation of its mechanism of action through scanning electron microscopy. *Microbial pathogenesis*, 136, 103716. <https://doi.org/10.1016/j.micpath.2019.103716>.
- [2]. Andrew AS, Burgess JL, Meza MM, Demidenko E, Waugh MG, Hamilton JW, Karagas MR. 2006. Arsenic exposure is associated with decreased DNA repair in vitro and in individuals exposed to drinking water arsenic. *Environ Health Perspect.* Aug;114(8):1193-8.
- [3]. Bennett MA, Huang TN, Matheson TW, Smith AK, Nickerson W. (η⁶- Hexamethylbenzene) ruthenium complexes. *Inorganic Syntheses* 1982; 21: 74–78. 58.
- [4]. Bones RW and Tausky HH. 1945. Colorimetric determination of creatinine by the Jaffe reaction.
- [5]. *J. Biol. Chem.* 158:581-591
- [6]. Carter DE, et al. 2003. The metabolism of inorganic arsenic oxides, gallium arsenide, and arsine: a toxicological review. *Toxicol Appl Pharmacol.*;193:309–34.
- [7]. Chen Y, Parvez F, Liu M, et al. 2011. Association between arsenic exposure from drinking water and proteinuria: results from the Health Effects of Arsenic Longitudinal Study. *Int J Epidemiol.*;828–35.
- [8]. Cohen SM, Arnold LL, Eldan M, Lewis AS, Beck BD. 2006. Methylated arsenicals: the implications of metabolism and carcinogenicity studies in rodents to human risk assessment. *Crit. Rev. Toxicol.*, 36: 99–133.
- [9]. Draper HH, Hadley M. 1990. Malondialdehyde determination as index of lipid peroxidation.
- [10]. *Methods Enzymol*, 186: 421-431.
- [11]. Duker AA, Carranza EJM, Hale M 2005. Arsenic geochemistry and health. *Environ. Int.* 31:631- 641.
- [12]. Dwivedi, N., & Flora, S. J. 2015. Sub-chronic exposure to arsenic and dichlorvos on erythrocyte antioxidant defense systems and lipid peroxidation in rats. *Journal of environmental biology*, 36(2), 383–391.
- [13]. Fawcett JK and Scott JE. 1960. A rapid and precise method for the determination of urea. *J. Chem. Pathol.* 13:156.

- [15]. Gallo M, Ferracane R, Giulia G, Ritieni A, Fagliano V. Microwave assisted extraction of phenolic compounds from four different spices. *Molecules* 2010; 15(3): 6366-6374.
- [16]. Guo S., Duan J. A., Tang Y. P., Qian Y. F., Zhao J. L., Qian D. W., Su S. L., Shang E. X., 2011.
- [17]. *J. Pharmaceut. Biomed.*,56(2), 264.
- [18]. Hajlaoui H, Mighri H, Noumi E, Snoussi M, Trabelsi N, Ksouri R. Chemical composition and biological activities of Tunisian *Cuminum cyminum* L. essential oil: A high effectiveness against *Vibrio* spp. strains. *Food Chem Toxicol* 2010;48: 2186-92.
- [19]. Hartwig A, Asmuss M, Ehleben I, Herzer U, Kostelac D, Pelzer A, Schwerdtle T, Bürkle. 2002. A. Interference by toxic metal ions with DNA repair processes and cell cycle control: molecular mechanisms. *Environ Health Perspect.*;110 Suppl 5:797-9.
- [20]. Hassan M 2018. Arsenic in groundwater. CRC Press, Boca Raton.
- [21]. Helleday T, Nilsson R, Jenssen D. 2000.Arsenic [III] and heavy metal ions induce intrachromosomal homologous recombination in the hprt gene V79 Chinese hamster cells. *Environ. Mol. Mutagen.*, 35: 114-122.
- [22]. Hsueh YM, Chung CJ, Shiue HS, et al. 2009. Urinary arsenic species and CKD in a Taiwanese population: a case-control study. *Am J Kidney Dis.* 54:859-70.
- [23]. Huang C, Ke Q, Costa M, Shi X. 2004. Molecular mechanisms of arsenic carcinogenesis. *Mol. Cell Biochem.* 255: 57-66.
- [24]. Jendrassik GF and Grofs BM. 1938. Quantitative colorimetric determination of bilirubin in serum or plasma. *Clin.Chem. Acta.* (27):79.
- [26]. Jha V, Garcia-Garcia G, Iseki K, et al., 2013.Chronic kidney disease: global dimension and perspectives. *Lancet.* 382:260-72.
- [27]. Johri RK. *Cuminum cyminum* and *Carum carvi*: An update. *Pharmacogn Rev* 2011; 5(9): 63-72.
- [28]. Kannan GM, Tripathi N, Dube SN and Flora SJS 2001. Toxic effects of arsenic (III) on some hematopoietic and central nervous system variables in rats and guinea pigs. *Journal of Toxicology. Clinical Toxicology* 39(7): 675-682.
- [29]. Kenneth GB, Gilbert LR. 2002. Arsenic, drinking water, and health: a position paper of the American council on science and health. *Regulatory Toxicology and Pharmacology*36(2): 162-174.
- [30]. Kind PRH and King EJ. 1954. Determination of Alkaline Phosphatase activity in serum. *J. Clin. Path.* 7: 322.
- [32]. Kumar A, Ali M, Kumar R, Rahman S. M, Srivastava A, Chayal K. N, Sagar V, Kumari R, et al. 2020. High Arsenic Concentration in Blood Samples of People of Village Gyaspur Mahaji, Patna, Bihar Drinking Arsenic-Contaminated Water. *Springer Nature Journal Exposure and Health*, 12, 131-140.
- [33]. Kumar, P., Kumar, M., Ramanathan, A. L., & Tsujimura, M. 2010. Tracing the factors responsible for arsenic enrichment in groundwater of the middle Gangetic Plain, India: a source identification perspective. *Environmental geochemistry and health*, 32(2), 129-146. <https://doi.org/10.1007/s10653-009-9270-5>.
- [34]. Li R, Jiang ZT. Chemical composition of the essential oil of *Cuminum cyminum* L. from China. *Flav Fragr J* 2004; 19: 311-313.
- [36]. Lu M, Wang H, Li XF, Arnold LL, Cohen SM, Le XC., 2007. Binding of dimethylarsinous acid to cysteine-13alpha of rat hemoglobin is responsible for the retention of arsenic in rat blood. *Chem Res Toxicol.*;20(1):27-37.
- [37]. Mallikarjuna GU, Dhanalakshmi S, Raisuddin S, Rao AR. 2003. Chemomodulatory influence of *Ferula assafoetida* on mammary epithelial differentiation, hepatic drug metabolizing enzymes, antioxidant

profiles and N-methyl-N-nitrosourea-induced mammary carcinogenesis in rats. *Breast Cancer Res Treat.*;81:1–10.

- [38]. Maurya, U. and Srivastava, S. 2011. Traditional Indian herbal medicine used as antipyretic, antiulcer, anti-diabetic and anticancer: A review. *International Journal of Research in Pharmaceutical Chemistry*, 4:1152-9..
- [39]. Miller, W.H.; Schipper, H.M.; Lee, J.S.; Waxman, S. 2002. Mechanisms of action of arsenic trioxide. *Cancer Res.*, 62: 3893-3903.
- [40]. Morshedi D, Aliakbari F, Tayaranian M, Fassihi P. 2015. Cuminaldehyde as the major component of cuminum cyminum, a natural aldehyde with inhibitory effect on alpha synuclein fibrillation and cytotoxicity. *Journal of Food Science*: 80(10): H2336-H2345.
- [41]. Morshedi D, Aliakbari F. 2012. The inhibitory effects of cuminaldehyde on amyloid fibrillation and cytotoxicity of alpha-synuclein. *Modares Journal of Medical Sciences: Pathobiology*; 15: 45-60.
- [42]. NIOSH (National Institute for Occupational Safety and Health) 1994). Elements in blood or tissue. Method: 8005. In: NIOSH manual of analytical methods. 4 Edn. O.H. Cincinnati: Department of Health and Human Service, National Institute of Occupational Safety and Health. *DHHS publication*; 94–113.
- [43]. Patlolla, A.K., Tchounwou, P.B., 2005. Serum acetyl cholinesterase as a biomarker of arsenic induced neurotoxicity in Sprague-Dawley rats. *International Journal of Environmental Research and Public Health*, 2: 80-83.
- [44]. Rahman MS, Kumar A, Kumar R, Ali M, Ghosh AK, Singh SK. Comparative Quantification Study of Arsenic in the Groundwater and Biological Samples of Simri Village of Buxar District, Bihar, India. *Indian J Occup Environ Med* 2019; 23:126–132.
- [45]. Reitman's S. and Frankel's S. 1957. A colorimetric method for determination of serum glutamate oxalacetic and glutamic pyruvate transaminases." *Amer J Clin Path* 1957; 28(1): 56-63.
- [46]. Saha D. 2009. Arsenic groundwater contamination in parts of middle Ganga plain, Bihar. *Current Science*. 97(6):753-755.
- [47]. Saha, D., & Sahu, S. 2016. A decade of investigations on groundwater arsenic contamination in Middle Ganga Plain, India. *Environmental geochemistry and health*, 38(2), 315–337. <https://doi.org/10.1007/s10653-015-9730-z>.
- [48]. Sankar, P., Telang, A. G., Kalaivanan, R., Karunakaran, V., Suresh, S., & Kesavan, M. 2016. Oral nanoparticulate curcumin combating arsenic-induced oxidative damage in kidney and brain of rats. *Toxicology and industrial health*, 32(3), 410–421. <https://doi.org/10.1177/0748233713498455>.
- [49]. Sekine T, Sugano M, Majid A, Fujii Y. 2007. Antifungal effects of volatile compounds from black zira (*Bunium persicum*) and other spices and herbs. *Journal of chemical ecology*; 33: 2123-32.
- [50]. Sharma, A., Kshetrimayum, C., Sadhu, H. G., & Kumar, S. 2018. Arsenic-induced oxidative stress, cholinesterase activity in the brain of Swiss albino mice, and its amelioration by antioxidants Vitamin E and Coenzyme Q10. *Environmental science and pollution research international*, 25(24), 23946–23953. <https://doi.org/10.1007/s11356-018-2398-z>.
- [51]. Singh RP, Gangadharappa HV, Mruthunjaya K. 2017. Cuminum cyminum – A Popular Spice: An Updated Review. *Pharmacogn J.*;9(3):292-301.
- [52]. Singh SK, Ghosh AK, Kumar A, Kislay K, Kumar C, Tiwari RR, et al. 2014. Groundwater arsenic contamination and associated health risks in Bihar, India, *Int J Environ Res*. 8:49.60.
- [53]. Smith AH, Goycolea M, Haque R and Biggs ML 1998. Marked increase in bladder and lung cancer mortality in a region of northern Chile due to arsenic in drinking water. *American Journal of Epidemiology* 147(7): 660–669.

- [54]. Soni KB, Rajan A, Kuttan R. 1993. Inhibition of aflatoxin-induced liver damage in ducklings by food additives. *Mycotoxin Res.*;9:22–7.
- [55]. Styblo M, et al. 2002. The role of biomethylation in toxicity and carcinogenicity of arsenic: a research update. *Environ Health Perspect.* 110(Suppl 5):767–71.
- [56]. Toro, G. et al. 1975. *practical clinical chem.*. P:154
- [57]. US NRC 2001. Arsenic in drinking water, 2001 update. Washington, DC, US National Research Council, National Academy Press.
- [58]. Vahidnia A, Romijn F, van der Voet GB and de Wolff F A 2008. Arsenic-induced neurotoxicity in relation to toxicokinetics: Effects on sciatic nerve proteins. *ChemicoBiological Interactions* 176(2-3): 188–195.
- [59]. Wang, J., Zhang, G., Lin, Z., Luo, Y., Fang, H., Yang, L., Xie, J., & Guo, L. 2020. Determination of arsenicals in mouse tissues after simulated exposure to arsenic from rice for sixteen weeks and the effects on histopathological features. *Ecotoxicology and environmental safety*, 200, 110742.
- [60]. Wanibuchi H, Yamamoto S, Chen H, Yoshida K, Endo G, Hori T, Fukushima S. 1996. Promoting effects of dimethylarsinic acid on N-butyl-N-(4-hydroxybutyl) nitrosamine-induced urinary bladder carcinogenesis in rats. *Carcinogenesis*.17:2435–2439.
- [61]. Wanner, J., Bail, S., Jirovetz, L., Buchbauer, G., Schmidt, E., Gochev, V., Girova, T., Atanasova, T., & Stoyanova, A. 2010. Chemical composition and antimicrobial activity of cumin oil (*Cuminum cyminum*, Apiaceae). *Natural product communications*, 5(9), 1355–1358.
- [62]. Yamamoto S, Konishi Y, Matsuda T, Murai T, Shibata MA, Matsui-Yuasa I, Otani S, Kuroda K, Endo G, Fukushima S. 1995. Cancer induction by an organic arsenic compound, dimethylarsinic acid (cacodylic acid), in F344/DuCrj rats after pretreatment with five carcinogens. *Cancer Res.*55:1271–1276.
- [63]. Yamanaka K, Kato K, Mizoi M, An Y, Takabayashi F, Nakano M, Hoshino M, Okada S. 2004.; The role of active arsenic species produced by metabolic reduction of dimethylarsinic acid in genotoxicity and tumorigenesis. *Toxicol. Appl. Pharmacol.*, 198: 385–393.
- [64]. Zhang R. T., Chen J. H., Shi Q., Li Z. Y., Peng Z. Y., Zheng L., Wang X. R., 2014. *Food Res. Int.*, 56(2), 47.
- [65]. Zheng L, Kuo CC, Fadrowski J, Agnew J, Weaver VM, Navas-Acien A. 2014. Arsenic and Chronic Kidney Disease: A Systematic Review. *Curr Environ Health Rep.* 1(3):192–207.
- [66]. Zheng LY, Umans JG, Tellez-Plaza M, et al. 2013. Urine arsenic and prevalent albuminuria: evidence from a population-based study. *Am J Kidney Dis.* 61:385–94.