

## Densities, Refractive Indices of Substituted Azomethine in Different Percent of Various Solvents

A.V.Kawalkar<sup>1</sup>, M. P. Wadekar<sup>2</sup>

<sup>1</sup>Department of Chemistry, Amolackchand Mahavidyalaya, Yavatmal, Maharashtra, India

<sup>2</sup>Department of Applied Chemistry Division, Govt. Vidarbha Institute of Science and Humanities, Amravati, Maharashtra, India

### ABSTRACT

Molecular interaction such as solute-solute, solute-solvent and solvent-solvent interactions in the substituted azomethine drug in the different percentage of organic solvent has been pointed out. In the present work refractive index and the densities of the substituted azomethine in different percent of various organic solvents were reported. The data thus helps to determine Molar refraction ( $R_m$ ) and polarizability constant ( $\alpha$ ) of some different substituted azomethine in binary mixture. Observations showed that the molar refraction and polarizability constant of substituted azomethine drugs increases with increase in percent composition of organic solvents.

**Keyword:** Substituted azomethine, molar refraction ( $R_m$ ), polarizability constant ( $\alpha$ ), refractometry, refractive index.

### I. INTRODUCTION

Refractive index of a liquid is very important property, which gives ideas about geometry and structure of molecule. The refractive index ( $n$ ) of the medium is the ratio of the velocity of light in vacuum to that in the medium. Its value depends upon the temperature and the wavelength of light used. Generally, the D-line of sodium is used for standard measurements. The refractive index is the ratio of angle of incident to the angle of refraction. Measurement of refractive index shows very interesting applications in pharmaceutical, chemical, agriculture, food, oil and beverage industries.

Many searchers have reported the refractive indices in mixed solvents [1-4]. The properties of liquid such as viscosity, refractive index and ultrasonic velocity of binary mixtures are studied by many workers [5-8]. Refractometric Study of S-trizinothiocarbamides in Dioxane-water was also reported. [9]. The viscometric, refractometric and interferometric measurements are very important in medicinal and drug chemistry role [10-12]. Oswal et al [13] have studied dielectric constants and refractive indices of binary mixtures. Dadhichi et al [14] have investigated the measurement of viscosity, refractivity index and metal ligand stability constant of substituted benzofurones in different solvents. Refractive indices of binary, ternary liquid solutions and solutions of biologically important compounds have been studied [15-21].

## II. MATERIAL & METHOD

In the present investigation, refractive indices of liquid mixtures were measured with the help of Abbe's refractometer, specially designed to measure the refractive indices of the small quantities of the transparent liquid by direct reading. The ligands of which physical parameters is to be explore are synthesized by using reported protocol.. The solutions of ligand in different percent composition of binary mixtures were prepared by weight. All the weighing were made on one pan digital balance (petit balance AD\_50B) with an accuracy of ( $\pm 0.001$ )gm.. The densities of solutions were determined by a precalibrated bicapillary pyknometer ( $\pm 0.1\%$ ). The constant temperature of the prism box is maintained by circulating water from thermostat at ( $27 \pm 0.1$ )OC.

### Calculation :

The molar refraction of solvent and solution are determined by using Lorentz- Lorentz equation.

The molar refraction of different solvent, mixtures are determined from-

$$R_{DMF-W} = X_1R_1 + X_2R_2 \quad (1)$$

where , R1 and R2 are molar refractions of DMF and water respectively.

The molar refraction of solutions of ligand in DMF-water mixtures are determined from-

$$\left( \frac{n^2 - 1}{n^2 + 2} \right) \left[ \frac{M_1}{d} + \frac{M_2}{d} + \frac{M_3}{d} \right] \dots \dots \dots (2)$$

where, n is the refractive index of solution, X1 is mole fraction of DMF, X2 is mole fraction of water and X3 is mole fraction of solute, M1, M2 and M3 are molecular weights of DMF, water and solute respectively. 'd' is the density of solution.

The molar refraction of ligand is calculated as –

$$R_{lig} = R_{mix} - R_{DMF} - w \quad (3)$$

The polarizability constant ( ) of ligand is calculated from following relation-

$$R_{lig} = \frac{4}{3} N_0 \dots \dots \dots (4) \text{ where, } N_0 \text{ is Avogadro's number.}$$

## III. RESULT AND DISCUSSION

Table 1: Values of Molar Refraction of different composition of solvents.

% of Solvent Mixture	Molar polarization Rm		
	DMSO	Dioxane	Ethanol
20%	15.0946	14.2357	19.1123
40%	14.2355	13.1155	18.5736
60%	12.7632	10.0759	15.0522
80%	10.8125	08.2301	12.7245
100%	5.7311	4.5711	7.0932
70%	10.2257	9.0325	14.0327

Table 2: The values of refractive index (n) and density(d) of 0.01M solution of ligand in different composition of DMSO, Dioxane and Ethanol solvent at 300K

% of Solvent Mixture	Molar polarization R <sub>m</sub>			Density (d) gm/cm <sup>3</sup>		
	DMSO	Dioxane	Ethanol	DMSO	Dioxane	Ethanol
Ligand L1						
20%	66.8917	57.6362	70.0866	1.0111	1.0051	1.0241
40%	78.0626	65.8380	81.6871	1.0187	1.0150	1.0277
60%	83.3022	70.8890	89.7373	1.0209	1.0166	1.0310
80%	91.9491	74.7821	95.5928	1.0255	1.0176	1.0339
100%	95.4783	79.5399	102.2347	1.0294	1.0213	1.0362
70%	84.7519	72.7507	92.1235	1.01923	1.01695	1.0321
Ligand L 2						
20%	79.5989	68.7954	86.9939	1.0063	1.0027	1.0072
40%	92.5314	78.7384	100.6731	1.0145	1.0131	1.0176
60%	99.3297	86.0426	109.0522	1.0181	1.0150	1.0237
80%	106.918	89.0521	117.4013	1.0291	1.0284	1.0278

100%	113.875	95.1013	123.66 69	1.029 7	1.0293	1.0313
70%	100.543	87.0155	114.10 21	1.019 5	1.0195	1.0257
Ligand L3						
20%	78.7969	67.8873	86.3531	1.0065	1.0042	1.0093
40%	92.1047	77.8228	101.464 8	1.0095	1.0087	1.0130
60%	99.6751	86.2908	109.898 7	1.0105	1.0090	1.0231
80%	105.515 2	90.0569	116.322 6	1.0205	1.0197	1.0267
100%	111.997 3	95.5152	121.780 5	1.0223	1.0203	1.0310
70%	101.212 3	88.3011	113.637 2	1.0168	1.0147	1.0253
Ligand 4						
20%	94.9043	82.6679	102.59 96	0.9999	0.9987	1.0119
40%	111.382 3	94.0953	120.67 84	1.0163	1.0150	1.0195
60%	119.320 5	102.837 9	130.24 51	1.0211	1.0199	1.0232
80%	126.449 6	107.741 9	138.76 19	1.0283	1.0256	1.0276

100%	133.291 6	113.156 8	146.55 56	1.0299	1.0257	1.0357
70%	123.001 6	104.450 9	134.24 15	1.0235	1.0221	1.0255

Table 3: The values of Molar refraction ( $R_m$ ), polarizability constant ( $\alpha$ ) of 0.01M solution of ligand in different composition of DMSO, Methanol and Acetone solvent at 300K.

% of Solvent Mixture	Molar refraction ( $R_m$ ) $\times 10^3 \text{ cm}^3/\text{mole}$			polarizability constant ( $\alpha$ ) $\times 10^{-23} \text{ cm}^3$		
Ligand L1						
	DMSO	Dioxane	Ethanol	DMSO	Dioxane	Ethanol
20%	66.8917	57.6362	70.0866	2.3237	2.2856	2.7794
40%	78.0626	65.8380	81.6871	2.6975	2.6109	3.2394
60%	83.3022	70.8890	89.7373	2.8732	2.8112	3.5587
80%	91.9491	74.7821	95.5928	2.9955	2.9657	3.7909
100%	95.4783	79.5399	102.2347	3.1945	3.1543	4.0543
70%	84.7519	72.7507	92.1235	2.9511	2.9019	3.6751
Ligand L2						
20%	79.5989	68.7954	86.9939	2.8327	2.7282	3.4499
40%	92.5314	78.7384	100.6731	3.1923	3.1225	3.9923

60%	99.3297	86.0426	109.0522	3.4812	3.4121	4.3246
80%	106.9179	89.0521	117.4013	3.5985	3.5315	4.6557
100%	113.8750	95.1013	123.6669	3.8618	3.7714	4.9042
70%	100.543	87.0155	114.1021	3.5534	3.4528	4.4836
Ligand L3						
20%	78.7969	67.8873	86.3531	2.7357	2.6922	3.4245
40%	92.1047	77.8228	101.4648	3.1845	3.0861	4.0237
60%	99.6751	86.2908	109.8987	3.5193	3.4220	4.3582
80%	105.5152	90.0569	116.3226	3.6327	3.5713	4.6129
100%	111.9973	95.5152	121.7805	3.8125	3.7878	4.8294
70%						
	101.2123	88.3011	113.6372	3.5579	3.4911	4.5109
Ligand L4						
20%	94.9043	82.6679	102.5996	3.3115	3.2783	4.0687
40%	111.3823	94.0953	120.6784	3.7542	3.7315	4.7875
60%	119.3205	102.8379	130.2451	4.1528	4.0782	5.1651

80%	126.449 6	107.7419	138.7619	4.3272	4.2727	5.5028
100%	133.291 6	113.1567	146.5556	4.5545	4.4874	5.8119
70%	123.001 6	104.4509	134.2415	4.2275	4.1543	5.3415

Graphical representation of molar polarization ( $R_m$ ) of all ligand at 0.01M verses concentration in different percentage of DMSO solvent







Fig. 1  $R_m$  verses concentration of ligand L1

Graphical representation of molar polarization ( $R_m$ ) of all ligand at 0.01M verses concentration in different percentage of 1,4 Dioxane solvent





Graphical representation of molar polarization ( $R_m$ ) of all ligand at 0.01M verses concentration in different percentage of Ethanol solvent



The value of molar refraction of different percent composition in binary mixture are shown in table-1. From the data it is observed that value of molar refraction goes on increasing with the decrease in amount of water in percent composition. Comparatively molar refraction of DMSO is greater than acetone and methanol this is due to more value of dipole moment of DMSO.

Table-2 shows the comparative data of refractive indices and densities of DMSO, acetone and methanol in different percent composition with water. From this, it is observe that, refractive index and density increases with the increase in percent composition of organic solvent. Graphical representation between molar refraction and percent composition of DMSO, methanol and acetone shows linear relationship. (Fig.1-5 DMSO, fig.6-10 methanol, fig.11- 15 acetone) Those solvent having more value of dipole moment shows greater refractive index and density, also there is same trend in case of ligand used. Ligand having more dipole moment shows greater value of refractive index and less value of density.

Table-3 shows the comparative data of molar refraction and polarizability constant. These parameter provide important information about structural orientation of ligand in solution. From this it is observed that, molar refraction and polarizability constant in methanol is higher than DMSO and acetone. The trend regarding

increasing value of molar refraction and polarizability constant is methanol > DMSO > acetone. From this observation it is concluded that, methanol has strong hydrogen bonding, which make solution more viscous which is responsible for more bending of light towards normal. In case of DMSO, it has more value of molar refraction and polarizability constant than acetone because it has more dipole moment.

#### IV. REFERENCES

- [1] . S D Devsarkar, V V Pandhare, P S Kattakar, J. Engineering, 2013, (2012), 1-4.
- [2] . S S Dhondge, J. Chem. Eng. Data, 55, (2010), 3962-3968.
- [3] . G B Pethe, A A Ramteke, T R Lawankar, J Chem. Pharm., 2(4), (2010), 68-74.
- [4] . S P Wagh, Ph.D Thesis in Chemistry, Amravati University, Amravati, (2004).
- [5] . D T tayade, A M Kshirsagar, The open physical chemistry Journal, 6, (2014), 1-7.
- [6] . G. satyamaiah, M Prasad, M chandrashekhar, Indian J. Advance in chemical science, 2(2),(2014), 116-123..
- [7] . M I Aralaguppi, T M Aminabhavi, R H Balundgi , S S Joshi, J Phys Chem, 95 ,(1991), 5299.
- [8] . A S Burghate, P B Agrawal, S W Quazi, M L Narwade, Asian J Chem, 13(4), (2001), 1652
- [9] . D. T. Tayade, Sheikh Oriental Journal Of Chemistry, 28(1), ( 2012), 559-564
- [10] . S Nagar., H Singh, J. Med. Chem., 16 ,(2007), 178-180.
- [11] . L. Hall, J. Phys. Rev.,(1998), 73-76.
- [12] . J. Pandey, et .al Shukla, J. Chem. Eng. Data, 34, ( 1989), 29-32.
- [13] . S L Ostwal , M V Rathnam. Indian J. Chem., 26 (1987), , 29.
- [14] . V S Jamode et.al J.Ind.Coun.of Chemist, 35 ,(2005),22,.
- [15] . M. V. Rathnam , S. Mohite Journal of the Serbian Chemical Society, 77(4), (2012). 507-521,
- [16] . A. Arce et.al, Journal of Solution Chemistry, vol. 27(10), (1998), 911-923,
- [17] . R. Talegaonkar et.al Oriental Journal Of Chemistry, 27(3) ,(2011).1285-1288,
- [18] . P. Baraldi et.al Journal of Solution Chemistry, 31(11), (2002)., 873-893,
- [19] . A. Arce et.al Journal of Chemical and Engineering Data, 42( 4), (1997). 721-726,
- [20] . A. N. Sonar , N. S. Pawar, " Rasayan Journal of Chemistry, 3( 2), (2010) 250-254.
- [21] . S. C. Bhatia et al International Journal of ermophysics, vol. 32( 6), (2011). 1163-1174,