

# An Overview on Oxidative Stress in Diabetes Mellitus

Jagroop Singh<sup>1</sup>, Dr. Sukhraj Kaur<sup>2</sup>, Dr. Manpreet Kaur Verma<sup>3</sup>, Manjinder Kaur<sup>4</sup>

<sup>1</sup>Ph.D (Biochemistry) Scholar, Desh Bhagat University, Mandi Gobindgarh, Punjab, India

<sup>2</sup>Professor and Head, Department of Biochemistry, Govt Medical College Amritsar Punjab, India

<sup>3</sup>Assistant Professor, Department of Chemistry, Desh Bhagat University, Gobindgarh, Punjab, India

<sup>4</sup>Nursing Officer, Government Medical college Amritsar Punjab, India

## ABSTRACT

Human body is continuously exposed to different types of agents that results in the production of reactive species called as free radicals (ROS/RNS) which by the transfer of their free unpaired electron causes the oxidation of cellular machinery. In order to encounter the deleterious effects of such species, body has got endogenous antioxidant systems or it obtains exogenous antioxidants from diet that neutralizes such species and keeps the homeostasis of body. Any imbalance between the RS and antioxidants leads to produce a condition known as “oxidative stress” that results in the development of pathological condition among which one is diabetes. Most of the studies reveal the inference of oxidative stress in diabetes pathogenesis by the alteration in enzymatic systems, lipid peroxidation, impaired Glutathione metabolism and decreased Vitamin C levels. Lipids, proteins, DNA damage, Glutathione, catalane and superoxide dismutase are various biomarkers of oxidative stress in diabetes mellitus. Oxidative stress induced complications of diabetes may include stroke, neuropathy, retinopathy and nephropathy. The basic aim of this review was to summarize the basics of oxidative stress in diabetes mellitus

**Keywords** : Diabetes Mellitus, Oxidative Stress, Biomarkers.

## Article Info

Volume 9, Issue 3

Page Number : 444-448

## Publication Issue

May-June-2022

## Article History

Accepted : 01 June 2022

Published : 07 June 2022

## I. INTRODUCTION

Diabetes mellitus, is a chronic metabolic disorder characterized by a high blood glucose concentration hyperglycemia (fasting plasma glucose > 7.0 mmol/l or plasma glucose > 11.1 mmol/l 2 hours after a meal) – caused by insulin deficiency, often combined with insulin resistance [1]. Diabetes mellitus is a complicated metabolic disorder that has gravely


troubled the human health and quality of life. It is a global health crisis, which has been persistently affecting the humanity, irrespective of the socioeconomic profile and geographic location of the population. According to an estimate, every five seconds one person is detected with diabetes somewhere in the world, while someone dies of it every ten seconds [2]. Oxidative stress plays a major role in the onset and progression of diabetes. The

development of oxidative stress in the body leads to the generation of free radicals which in turn causes major damage to the body [3]. Further the onset of diabetes is promoted by the oxidative stress as it causes damage to the  $\beta$ -cells and reduces insulin sensitivity [4]. Most common risk factors includes as increased age, unhealthy life style and obesity, all contribute to an oxidative environment that may increase insulin resistance or impairing glucose tolerance. The process by which alteration in insulin sensitivity occurs is often multifactorial and quite complex, involving multiple cell signaling pathways. A common outcome of diabetes is hyperglycemia, which in turn contributes to the progression and disturbance of an overall cellular oxidative status [5]. This review discusses the role of oxidative stress in the diabetes process.

## II. Oxidative Stress And Diabetes

The human body is incessantly exposed to free radicals from inside the body (endogenous) and outside the body (exogenous). For production of energy cells need continuous supply of oxygen. During the process of metabolic respiration, the cells take in oxygen, burn it, and release energy [6]. Throughout these metabolic processes, free radicals species are generated; however these free radicals are scavenged by the body's internal enzymatic and non-enzymatic antioxidant army. Oxidative stress occurs when there is an imbalance occurs between free radical production and scavenging by the antioxidants. This imbalance happens for one of two reasons: a) when the body's antioxidant status is decreased, or b) when excessive free radical species are generated [6]. For instance in diabetes, due to hyperglycemia there is an increased speed of endogenous free radical generation and reduction in antioxidant status, which leads to diabetic oxidative stress. It is this oxidative stress which plays pivotal role in the progression and the development of diabetes and its complications [6]. Ha et al (2000),

reported that oxidative stress is one of the prominent factor for vascular complications in diabetes including nephropathy [6]. Several mechanisms, including autoxidative glycosylation, formation of AGEs (advanced glycation end products), and increased polyol pathway activity contribute to increased oxidative stress as shown in Figure 1. In hyperglycemic condition auto-oxidation of glucose can generate tremendous reactive oxygen species, which is believed to be the main source of free radicals. During hyperglycemic condition glucose in presence of transition metals gets oxidized to an enediol radical anion that is converted into superoxide radical species and ketoaldehydes. The superoxide anion radicals gets dismutated into hydrogen peroxide, which in the presence of transitional metals can lead to production of extremely reactive hydroxyl radicals [7]. Superoxide anion radicals can also react with nitric oxide to form reactive peroxynitrite radicals. During hyperglycemia, lipid peroxidation of low density lipoprotein (LDL) can be promote by a superoxide dependent pathway to produce free radical species [8].



**Fig 1. Hyperglycaemia induced oxidative Stress**

Another important free radical generating source in diabetes is protein glycation. In glycation condensation of glucose with the  $\epsilon$ -amino group of lysine, the  $\alpha$ -amino group of an N-terminal amino acid or the amines of nucleic acids occurs, which leads to the formation of advanced glycation end products (AGEs). Glycation has been the major source of

increased generation of ROS in diabetes patients [9]. AGEs are supposed to be responsible in the genesis of most of the irreversible complications of diabetes, including vascular complication expanded extracellular matrix, hyperplasia, and cellular hypertrophy [10]. In addition to glucose induced oxidative stress by glycol-oxidation product formation, fructose which is elevated as a result of polyol pathway activation also leads to the formation of AGE precursor's like glyoxyl, methylglyoxal and 3-deoxyglucosone [11]. These AGEs, via their receptors (RAGEs), inactivate enzymes by altering their structures, increases free radicals formation, and suppress the activity of nitric oxide. Elevation in intracellular oxidative stress results in activation of the transcription factor NF- $\kappa$ B, which promotes up-regulation of various NF- $\kappa$ B controlled target genes [12]. NF- $\kappa$ B promotes generation of nitric oxide, which is supposed to be a mediator of pancreatic  $\beta$ -cell damage. In addition, prolonged hyperglycemia promotes glycation of antioxidant enzymes, which could alter their structure and function such that they are unable to scavenge free radicals, exacerbating oxidative stress in diabetes [12]. Therefore, the process of glucose oxidation promotes not only increased ROS products but also for decrease bioavailability of cellular antioxidant status.

Biomarkers of Diabetic oxidative stress;

Antioxidant defense system in the cell includes glutathione (GSH), as well as antioxidant enzymes such as catalase (CAT), superoxide dismutase (SOD), glutathione peroxidase (GPx), glutathione S-transferase (GST), and glutathione reductase (GRx). Because of their ability to scavenge for and attenuate the effects of ROS, these enzymes serve as biomarkers of oxidative stress.

Lipid peroxidation Induction of diabetes in rats with diabetogens like alloxan and streptozotocin uniformly promotes an increase in thiobarbituric acid reactive substances (TBARS), an indirect confirmation of intensified freeradical production. MDA has been

considered as a potential primary biomarker of free radical mediated lipid damage and oxidative stress [13]. Elevation in lipid peroxidation is also a confirmation of decline in defense mechanisms of cellular antioxidant currency [14]. Lipid peroxidation of cellular structures promotes atherosclerosis in diabetic patients. In diabetic patients, TBARS in the red blood cells as well as in serum are significantly higher while as reduction in erythrocyte antioxidant status [15, 16]. Thus preventing/reducing the formation of reactive oxygen species would be an efficient means to reduce free radical induced cellular damage, and several antioxidant compounds have been tested in diabetic animal models with varying success.

#### **Glutathione Levels**

Reduced glutathione is an important cellular redox buffer having concentrations up to 10 mM [17]. Glutathione may act as a direct free-radical scavenger, as a cofactor for many enzymes, and as a co-substrate for glutathione peroxidase activity [18]. In chemically induced diabetic animals glutathione concentration is found to be decreased in the kidney [19], liver [16] pancreas [20]. Glutathione Peroxidase and Glutathione Reductase Glutathione peroxidase (GPx) converts harmful hydrogen peroxide to water in presence of reduced glutathione as a co-substrate [21]. Glutathione reductase converts oxidized glutathione back to glutathione using the NADPH as a cofactor. In chemically induced diabetic animals glutathione peroxidase and glutathione reductase concentration is found to be decreased in liver [22] and pancreatic [20] tissue.

#### **Superoxide Dismutase**

Superoxide dismutase (SOD) is one of the important antioxidant enzyme that neutralize the highly reactive oxygen species superoxide anion ( $O_2^-$ ) by dismutation into hydrogen peroxide and molecular oxygen [23]. SOD protects against cellular and histological damages that are produced by ROS which

in turn reduces diabetic complications [24]. During development of diabetic nephropathy, superoxide dismutase acts as a major defender in its progression. Down regulation of renal SOD promotes the progression of diabetic nephropathy [25].

### Catalase

Nearly every living organism is bestowed by catalase due its potential antioxidant activity. It has been reported that catalase plays a vital role in attenuating oxidative stress-generated diabetic complications [26]. During oxidative stress pancreatic cells are protected by catalase from destructive hydrogen peroxide reactive species [27].

### III.CONCLUSION

Oxidative stress has been demonstrated in many studies to participate in the progression of diabetes which plays important role during diabetes, including impairment of insulin action and elevation of the complication incidence. Antioxidants have already shown to be prospective in the treatment of diabetes both type 1 and type 2. Increase in the levels of oxygen and nitrogen free radicals (ROS/RNS) has been linked with lipid peroxidation, non-enzymatic glycation of proteins and oxidation of glucose which contributes toward diabetes mellitus and its complications. Most of the studies have shown relationship between oxidative stress and diabetes along with their complications related to heart, liver kidney and eye. Thus, oxidative stress seems to be more worrying in metabolic disorders specially diabetes type 2.

### IV. REFERENCES

- [1]. Prasad, S.K., Kulshreshtha, A., and Qureshi, T.N. (2009). Antidiabetic activity of some herbal plants in streptozotocin induced diabetic albino rats. *Pak J Nutr* 8, 551-557.
- [2]. Colagiuri, R. (2010). Diabetes: a pandemic, a development issue or both? *Expert review of cardiovascular therapy* 8, 305.
- [3]. Ali, A., Dar, M. A., & Ayaz, A. (2017). Diagnostic Approaches to Diabetes Mellitus and the Role of Vitamins. *J Nutr Food Sci*, 7(601), 2.
- [4]. Ali, A., Dar, M. A., & Ayaz, A. (2017). Diagnostic Approaches to Diabetes Mellitus and the Role of Vitamins. *J Nutr Food Sci*, 7(601), 2.
- [5]. K, A.F., H, Z., and L, K.G. (2005). Molecular targets of diabetic cardiovascular complications. *Curr. Drug Targets* 6, 487-494.
- [6]. Ha, H., and Lee, H.B. (2000). Reactive oxygen species as glucose signaling molecules in mesangial cells cultured under high glucose. *Kidney international* 58, S19-S25.
- [7]. Jiang, Z.-Y., Woollard, A.C.S., and Wolff, S.P. (1990). Hydrogen peroxide production during experimental protein glycation. *FEBS letters* 268, 69-71.
- [8]. Tsai, E.C., Hirsch, I.B., Brunzell, J.D., and Chait, A. (1994). Reduced plasma peroxy radical trapping capacity and increased susceptibility of LDL to oxidation in poorly controlled IDDM. *Diabetes* 43, 1010-1014.
- [9]. Wolff, S.P. (1993). Diabetes mellitus and free radicals Free radicals, transition metals and oxidative stress in the aetiology of diabetes mellitus and complications. *British medical bulletin* 49, 642-652.
- [10]. Munch, G., Thome, J., Foley, P., Schinzel, R., and Riederer, P. (1997). Advanced glycation endproducts in ageing and Alzheimer's disease. *Brain research reviews* 23, 134-143.
- [11]. Takagi, Y., Kashiwagi, A., Tanaka, Y., Asahina, T., Kikkawa, R., and Shigeta, Y. (1995). Significance of fructose-induced protein oxidation and formation of advanced glycation end product. *Journal of diabetes and its complications* 9, 87-91.
- [12]. Mohamed, A.K., Bierhaus, A., Schiekofer, S., Tritschler, H., Ziegler, R., and Nawroth, P.P. (1999). The role of oxidative stress and NF- $\kappa$ B activation in late diabetic complications. *Biofactors* 10, 157-167.
- [13]. Williamson, J.R., Chang, K., Frangos, M., Hasan, K.S., Ido, Y., Kawamura, T., Nyengaard, J.R., van

- Den Enden, M., Kilo, C., and Tilton, R.G. (1993). Hyperglycemic pseudohypoxia and diabetic complications. *Diabetes* 42, 801-813.
- [14]. Ramalho, J.S., Marques, C., Pereira, P.C., and Mota, M.C. (1995). Role of glycation in human lens protein structure change. *European journal of ophthalmology* 6, 155-161.
- [15]. Mizisin, A.P., Kalichman, M.W., Bache, M., Dines, K.C., and DiStefano, P.S. (1998). NT-3 attenuates functional and structural disorders in sensory nerves of galactose-fed rats. *Journal of Neuropathology & Experimental Neurology* 57, 803-813.
- [16]. Thompson, K.H., and McNeill, J.H. (1993). Effect of vanadyl sulfate feeding on susceptibility to peroxidative change in diabetic rats. *Research communications in chemical pathology and pharmacology* 80, 187-200.
- [17]. Griffith, O.W. (1980). Determination of glutathione and glutathione disulfide using glutathione reductase and 2-vinylpyridine. *Analytical biochemistry* 106, 207- 212.
- [18]. Josephy, P.D. (1997). *Molecular Toxicology*. New York: Oxford University Press.
- [19]. Aragno, M., Tamagno, E., Gatto, V., Brignardello, E., Parola, S., Danni, O., and Boccuzzi, G. (1999). Dehydroepiandrosterone protects tissues of streptozotocin-treated rats against oxidative stress. *Free Radical Biology and Medicine* 26, 1467-1474.
- [20]. Abdel-Wahab, M.H., and Abd-Allah, A.R. (2000). Possible protective effect of melatonin and/or desferrioxamine against streptozotocin-induced hyperglycaemia in mice. *Pharmacological Research* 41, 533-537.
- [21]. Sies, H. (1993). Damage to plasmid DNA by singlet oxygen and its protection. *Mutation Research/Genetic Toxicology* 299, 183-191.
- [22]. El-Missiry, M.A., and El Gindy, A.M. (2000). Amelioration of alloxan induced diabetes mellitus and oxidative stress in rats by oil of *Eruca sativa* seeds. *Annals of Nutrition and Metabolism* 44, 97-100.
- [23]. Kim, S.-S., Gallaher, D.D., and Csallany, A.S. (2000). Vitamin E and probucol reduce urinary lipophilic aldehydes and renal enlargement in streptozotocin-induced diabetic rats. *Lipids* 35, 1225-1237.
- [24]. Ford, I., Cotter, M.A., Cameron, N.E., and Greaves, M. (2001). The effects of treatment with alpha]-lipoic acid or evening primrose oil on vascular hemostatic and lipid risk factors, blood flow, and peripheral nerve conduction in the streptozotocin-diabetic rat. *Metabolism* 50, 868-875.
- [25]. Santini, S.A., Marra, G., Giardina, B., Cotroneo, P., Mordente, A., Martorana, G.E., Manto, A., and Ghirlanda, G. (1997). Defective plasma antioxidant defenses and enhanced susceptibility to lipid peroxidation in uncomplicated IDDM. *Diabetes* 46, 1853-1858.
- [26]. Kedziora-Kornatowska KZ, Luciak M, Blaszczyk J, and W, P. (1998). Effect of aminoguanidine on erythrocyte lipid peroxidation and activities of antioxidant enzymes in experimental diabetes. *Clin Chem Lab Med* 36, 771-775.
- [27]. Van Dam, P.S., Van Asbeck, B.S., Van Oirschot, J., Biessels, G.J., Hamers, F.P.T., and Marx, J.J.M. (2001). Glutathione and lipoate in diabetic rats: nerve function, blood flow and oxidative state. *European journal of clinical investigation* 31, 417-424.

**Cite this article as :**

Jagroop Singh, Dr. Sukhraj Kaur, Dr. Manpreet Kaur Verma, Manjinder Kaur, "An Overview on Oxidative Stress in Diabetes Mellitus", *International Journal of Scientific Research in Science and Technology (IJSRST)*, Online ISSN : 2395-602X, Print ISSN : 2395-6011, Volume 9 Issue 3, pp. 444-448, May-June 2022. Available at doi : <https://doi.org/10.32628/IJSRST229393>  
Journal URL : <https://ijsrst.com/IJSRST229393>